

Incidence of q -Statistics in Rank Distributions

G. Cigdem Yalcin
Alberto Robledo
Murray Gell-Mann

SFI WORKING PAPER: 2014-07-024

SFI Working Papers contain accounts of scientific work of the author(s) and do not necessarily represent the views of the Santa Fe Institute. We accept papers intended for publication in peer-reviewed journals or proceedings volumes, but not papers that have already appeared in print. Except for papers by our external faculty, papers must be based on work done at SFI, inspired by an invited visit to or collaboration at SFI, or funded by an SFI grant.

©NOTICE: This working paper is included by permission of the contributing author(s) as a means to ensure timely distribution of the scholarly and technical work on a non-commercial basis. Copyright and all rights therein are maintained by the author(s). It is understood that all persons copying this information will adhere to the terms and constraints invoked by each author's copyright. These works may be reposted only with the explicit permission of the copyright holder.

www.santafe.edu



SANTA FE INSTITUTE

Incidence of q -statistics in rank distributions

G.Cigdem Yalcin ^{*}, Alberto Robledo [†], and Murray Gell-Mann [‡]

^{*}Department of Physics, Istanbul University, 34134, Vezneciler, Istanbul, Turkey, [†]Instituto de Fisica, y Centro de Ciencias de la Complejidad, Universidad Nacional Autonoma de Mexico, Apartado Postal 20-364, Mexico 01000 Distrito Federal, Mexico, and [‡]Santa Fe Institute, Santa Fe, NM 87501

Submitted to Proceedings of the National Academy of Sciences of the United States of America

We show that size-rank distributions with power-law decay (often only over a limited extent) observed in a vast number of instances in a widespread family of systems obey Tsallis statistics. The theoretical framework for these distributions is analogous to that of a nonlinear iterated map near a tangent bifurcation for which the Lyapunov exponent is negligible or vanishes. The relevant statistical-mechanical expressions associated with these distributions are derived from a maximum entropy principle with the use of two different constraints, and the resulting duality of entropy indexes is seen to portray physically relevant information. While the value of the index α fixes the distribution's power-law exponent, that for the dual index $2 - \alpha$ ensures the extensivity of the deformed entropy.

size-rank distributions | Tsallis statistics | duality of entropy

Zipf's law refers to the (approximate) power law obeyed by sets of data when these are sorted out and displayed by rank in relation to magnitude or rate of recurrence [1]. The sets of data originate from many different fields: astrophysical, geophysical, ecological, biological, technological, financial, urban, social, etc., suggesting some kind of universality. Over the years this circumstance has attracted much attention and the rationalization of this empirical law has become a common endeavor in the study of complex systems [2, 3]. Here we pursue further the view [4, 5] that an understanding of the omnipresence of this type of rank distribution hints to an underlying structure similar to that which confers systems with many degrees of freedom the familiar macroscopic properties described by thermodynamics. That is, the quantities employed in describing this empirical law obey expressions derived from principles akin to a statistical-mechanical formalism [4, 5]. The most salient result presented here is that the reproduction of the data via a maximum entropy principle indicates that access to its configurational space is severely hindered to a point that the allowed configurational space has a vanishing measure. This feature appears to be responsible for the entropy expression not to be of the Boltzmann-Gibbs or Shannon type but instead it takes that of the Tsallis form [6], while the extensivity of entropy is preserved. It is perhaps worth clarifying that our study is set in discrete space and it does not consider any formal Hamiltonian system.

In Fig. 1 we show three examples of ranked data that appear to display power-law behavior along a considerable large interval of rank values. In the top panels of this figure we show data for the wealth of billionaires in the US [7], in the middle panels data for the energy released by earthquakes in California [8], and in the bottom panels data for the intensity of solar flares [9]. In the left panels logarithmic scales are used for both size and rank, whereas the right panels show the same data in log-linear scales. The left panels indicate approximate power law decay for large rank and a clear deviation from this for small to moderate rank. As we shall show below the theoretical description reproduces the data in Fig. 1 for the entire rank interval.

In Section II we recall [4, 5, 10] the concise stochastic approach for raw data generated by a power-law distribution P for the size random variable N that yields an analytical expression for the size-rank distribution $N(k)$. This analytical expression involves a deformed exponential that has been

shown to reproduce quantitatively real data and has as a limiting form the classical Zipf law [4, 5, 10]. We also recall [4, 5] the analogy that exists between the stochastic approach and the deterministic nonlinear dynamics at and close to the tangent bifurcation. This analogy allows for a convenient description of finite-sized data that deviates from power-law behavior for both small and large rank. In Section III we derive the rank distribution $N(k)$ from a maximum entropy principle (MEP) and this allows us, via a well-known deformation index duality, to discuss two different entropy expressions of the Tsallis type obtained from two different sets of constraints [11, 12, 13]. The values of the two entropy expressions coincide but they yield different information for the set of data under consideration. In Section IV we use this duality to discuss entropy extensivity of the ranked data and the presence of a strong phase-space contraction. This is shown to be the source of a generalized entropy that departs from the usual Shannon expression. This departure is extreme for the classical Zipf case, implying that the data can sample only a set of zero measure. Finally, in Section V we discuss and summarize our results.

The distribution functions that generate Zipf's law

A basic approach for the study of ranked data consists of three simply related distribution functions [4, 5, 10]. The input is the distribution $P(N)$ of the data N under consideration, that is, it is assumed that the data is generated by a source described by $P(N)$ such that N can be thought of as a random variable. With no loss of generality we restrict N to take positive values within an interval $N_{\min} \leq N \leq N_{\max}$, where we allow for the limiting possibilities $N_{\min} = 0$ and/or $N_{\max} \rightarrow \infty$.

Significance

The contents presented are of prime importance to the field of generalized statistical mechanics. We fulfill a longstanding need of exhibiting the kind of abundant real world data that matches the formal developments in this subject. These are size-rank distributions for which we provide a solid bridge between experimental data and theory. Also, this work delivers a working explanation for the existing duality between the two Tsallis-type entropy expressions that generalize the canonical expression. One relates to the distributions power-law exponent whereas the other ensures entropy extensivity. The generalized entropies arise from a drastic reduction of configurations available to the system. We argue that this phase-space contraction is farthest for ranked data of the Zipf type.

Reserved for Publication Footnotes

The total number of data extracted from $P(N)$ is denoted by \mathcal{N} . Next, the (complementary) cummulative distribution $\Pi(N, N_{\max})$ is determined from $P(N)$,

$$\Pi(N, N_{\max}) = \int_N^{N_{\max}} P(N') dN', \quad [1]$$

where the normalization of $P(N)$ implies $\Pi(N_{\min}, N_{\max}) = 1$. We can recover $P(N)$ from $\Pi(N, N_{\max})$,

$$P(N) = -\frac{\partial}{\partial N} \Pi(N, N_{\max}). \quad [2]$$

By construction, the distribution $\Pi(N, N_{\max})$ sorts out data according to its magnitude: As N is decreased from N_{\max} the distribution Π increases monotonically taking values from $\Pi(N_{\max}, N_{\max}) = 0$ to $\Pi(N_{\min}, N_{\max}) = 1$, so it can be identified with k/\mathcal{N} , where k is the rank and \mathcal{N} is the total number of data extracted from $P(N)$, and $k_{\max} = \mathcal{N}$. The last and third distribution is the size-rank function $N(k)$ and can be obtained by solving

$$\frac{k}{\mathcal{N}} = \int_{N(k)}^{N_{\max}} P(N') dN', \quad [3]$$

for $N(k)$. If k is to be an integer the possible lower limits in the integral in Eq. (3), $N(1), N(2), \dots, N(k_{\max})$ are such that the integral takes values $1/\mathcal{N}, 2/\mathcal{N}, \dots, k_{\max}/\mathcal{N}$.

If we make use of a power law form for $P(N)$,

$$P(N) \sim N^{-\alpha}, \quad 1 \leq \alpha < \infty, \quad [4]$$

we have [10, 4, 5]

$$\begin{aligned} \Pi(N(k), N_{\max}) &= \int_{N(k)}^{N_{\max}} N^{-\alpha} dN \\ &= \frac{1}{1-\alpha} [N_{\max}^{1-\alpha} - N(k)^{1-\alpha}], \end{aligned} \quad [5]$$

or, in terms of the q -deformed logarithmic function $\ln_q(x) \equiv (1-q)^{-1}[x^{1-q} - 1]$ with q a real number,

$$\ln_{\alpha} N(k) = \ln_{\alpha} N_{\max} - \mathcal{N}^{-1} k. \quad [6]$$

The size-rank distribution $N(k)$ is explicitly obtained from the above with use of the inverse of $\ln_q(x)$, the q -deformed exponential function $\exp_q(x) \equiv [1 + (1-q)x]^{1/(1-q)}$, this is

$$N(k) = N_{\max} \exp_{\alpha}(-N_{\max}^{\alpha-1} \mathcal{N}^{-1} k). \quad [7]$$

When $\alpha = 1$ Eq. (7) acquires the ordinary exponential form

$$N(k) = N_{\max} \exp(-\mathcal{N}^{-1} k), \quad [8]$$

whereas in the limit $N_{\max} \rightarrow \infty$ Eq. (7) becomes the power law $N(k) \sim k^{1/(1-\alpha)}$ that when $\alpha = 2$ gives the simple classical Zipf's law form $N(k) \sim k^{-1}$.

An explicit analogy between the generalized law of Zipf and the nonlinear dynamics of intermittency has been studied [4, 5]. We recall the renormalization group (RG) fixed-point map for the tangent bifurcation. The trajectories x_t , $t = 1, 2, 3, \dots$, produced by this map, comply (analytically) with:

$$\ln_z x_t = \ln_z x_0 + ut \quad [9]$$

or

$$x_t = x_0 \exp_z [x_0^{z-1} ut] \quad [10]$$

where the x_0 are the initial positions. The parallels between Eqs. (9) and (10) with Eqs. (6) and (7), respectively, is clear, and therefore, we conclude that the dynamical system represented by the fixed-point map operates in accordance to the same q -generalized statistical-mechanical properties discussed below. We notice that the absence of an upper bound for the rank k in Eqs. (6) and (7) is equivalent to the tangency condition in the map. Accordingly, to describe data with finite maximum rank, we look at the changes in $N(k)$ brought about by shifting the corresponding map from tangency, *i.e.*, we consider the trajectories, x_t , with initial positions x_0 of the map:

$$x' = x \exp_z(ux^{z-1}) + \varepsilon, \quad 0 < \varepsilon \ll 1 \quad [11]$$

with the identifications $k = t$, $\mathcal{N}^{-1} = -u$, $N(k) = x_t + x^*$, $N_{\max} = x_0 + x^*$ and $\alpha = z$, where the translation, x^* , ensures that all $N(k) \geq 0$. The capability of this approach to reproduce quantitatively real data for ranked data with deviations from power law for large rank has been discussed [4, 5].

Rank distributions from maximum entropy principle

The rank distribution $N(k)$ described in the previous section can be obtained from a maximum entropy principle (MEP), and, as we shall see, this allows to put forward important interpretations regarding the nature of the systems that give rise to it. But first we adjust our interpretation of $N(k)$. This quantity is actually the size or magnitude of the data under consideration, the number of units that, in a microcanonical ensemble description, is the number of configurations that take place for a fixed value of k . Therefore its inverse, $p_k = 1/N(k)$, is the (uniform) probability for the occurrence of each unit that constitutes $N(k)$. The probability p_k is normalized for fixed k , and we denote its limiting values by $p_{\min} = 1/N_{\max}$ and $p_{\max} = 1/N_{\min}$, $N_{\min} \leq N(k) \leq N_{\max}$.

A formal investigation of the possible entropy expressions that generalize the Boltzmann-Gibbs or Shannon canonical form has been systematically carried out with the use of the MEP under the assumption that only three of the Shannon-Kinchin axioms hold [11, 12, 13]. (Inclusion of the fourth, composability, uniquely defines the canonical form). Here we focus only on the Tsallis expressions [14].

Consider the entropy functional $\Phi_1[p_k]$ with Lagrange multipliers a and b ,

$$\Phi_1[p_k] = S_1[p_k] + a \left[\sum_{k=0}^{k_{\max}} p_k - \mathcal{P} \right] + b \left[\sum_{k=0}^{k_{\max}} k p_k - \mathcal{K} \right], \quad [12]$$

where the entropy expression $S_1[p_k]$ has the trace form [11]

$$S_1[p_k] = \sum_{k=0}^{k_{\max}} s_1(p_k). \quad [13]$$

Optimization via $\partial \Phi_1[p_k] / \partial p_k = 0$, $k = 0, 1, 2, \dots, k_{\max}$, gives

$$s'_1(p_k) = -a - bk. \quad [14]$$

Now, the choices

$$s'_1(p_k) = \alpha \ln_{\alpha} p_k^{-1} - 1, \quad a = -\alpha \ln_{\alpha} p_{\min}^{-1} + 1, \quad b = \alpha \mathcal{N}^{-1}, \quad [15]$$

lead to

$$\ln_{\alpha} p_k^{-1} = \ln_{\alpha} p_{\min}^{-1} - \mathcal{N}^{-1} k. \quad [16]$$

or

$$p_k^{-1} = p_{\min}^{-1} \exp_{\alpha}(-p_{\min}^{1-\alpha} \mathcal{N}^{-1} k). \quad [17]$$

from which we immediately recover Eqs. (6) and (7).

We repeat the same optimization procedure but with a constraint change [11]. Consider the functional $\Phi_2[p_k]$ with Lagrange multipliers c and d ,

$$\Phi_2[p_k] = S_2[p_k] + c \left[\sum_{k=0}^{k_{\max}} p_k - \mathcal{P} \right] + d \left[\sum_{k=0}^{k_{\max}} k p_k^{\alpha'} - \mathcal{K}_{\alpha'} \right], \quad [18]$$

and where the entropy expression $S_2[p_k]$ has also a trace form

$$S_2[p_k] = \sum_{k=0}^{k_{\max}} s_2(p_k). \quad [19]$$

Optimization via $\partial \Phi_2[p_k] / \partial p_k = 0$, $k = 0, 1, 2, \dots, k_{\max}$, gives

$$s'_2(p_k) = -c - dk. \quad [20]$$

And this time the choices

$$s'_2(p_k) = -(2 - \alpha') \ln_{\alpha'} p_k - 1, \quad c = (2 - \alpha') \ln_{\alpha'} p_{\min} + 1, \quad [21]$$

$$d = (2 - \alpha') \mathcal{N}^{-1}, \quad [22]$$

give the expressions

$$\ln_{\alpha'} p_k = \ln_{\alpha'} p_{\min} + \mathcal{N}^{-1} k. \quad [23]$$

or

$$p_k = p_{\min} \exp_{\alpha'}(p_{\min}^{\alpha'-1} \mathcal{N}^{-1} k). \quad [24]$$

A comparison of Eqs. (6) and (7) with Eqs. (23) and (24), respectively, indicates that they become equivalent with the identifications

$$p_k = 1/N(k), \quad p_{\min} = 1/N_{\max}, \quad \alpha' = 2 - \alpha.$$

Furthermore, $s'_2(p_k) = s'_1(p_k)$ (as given by Eqs. (14), (15), (20) and (22)) and therefore

$$S_2[p_k] = S_1[p_k], \quad [25]$$

where their optimized expressions are

$$S_1[p_k] = \sum_{k=0}^{k_{\max}} p_k \ln_{\alpha} p_k^{-1}, \quad [26]$$

and

$$S_2[p_k] = - \sum_{k=0}^{k_{\max}} p_k \ln_{\alpha'} p_k. \quad [27]$$

Under the assumption of validity of only the first three Shannon-Kinchin axioms it has been shown [11, 12] that there are only two ways to construct entropy expressions via the MEP procedure. These correspond to the constraints used in Eqs. (12) and (18) and the resulting entropy expressions are those in Eqs. (26) and (27). The two approaches are related via the deformation index duality $\alpha' = 2 - \alpha$, and, for the same distribution p_k , their values are equal as in Eq. (25). For an earlier account of this duality property see Ref. [15]. See also [16]. From our earlier discussion we know that the index α fixes the shape of the rank distribution $N(k)$ and that its departure from unity generates its power-law feature and that the value $\alpha = 2$ reproduces the classic Zipf law. To complete the picture we need to clarify the role of the dual index α' and the distribution p_k , and from this obtain an understanding of the dual entropy expressions in Eqs. (26) and (27). Interestingly, when $\alpha = \alpha' = 1$ the duality collapses into

the Boltzmann-Gibbs or Shannon entropy expressions and the exponential form for $N(k)$, but for $\alpha = 2$ we have $\alpha' = 0$ and p_k grows linearly with k .

In Fig. 2 we show the same three sets of data in Fig. 1 in log-linear scales. This time we fit them with Eqs. (7) and (24) and observe that the data are well described with values of the deformations $\alpha = 2$ and $\alpha' = 0$.

Statistical mechanics of contracted configuration space

The function $N(k)$ has the properties of a microcanonical partition function [4, 5]. That is, the size $N(k)$ is the result of $N(k)$ equally-probable configurations, and the probabilities p_k are correspondingly normalized for fixed k . However, these probabilities are not normalized if the rank k runs across its values $k = 0, \dots, k_{\max}$, and we do not make an attempt here to do so. Instead, we look at the rank dependence Eq. (24), that we identify as the system's size dependence. As it can be observed in the right panels of Fig. 2 the probabilities p_k rises sharply and then saturates as k increases. The pure deformed exponential

$$\frac{p_k}{p_{\min}} = \frac{N_{\max}}{N(k)} = \exp_{\alpha'}(p_{\min}^{\alpha'-1} \mathcal{N}^{-1} k) \quad [28]$$

measures the change in the number of microcanonical configurations with the size of the system k . We define the size-dependent entropy

$$S(k) \equiv \ln_{\alpha'} \left(\frac{N_{\max}}{N_k} \right), \quad k \text{ fixed}, \quad [29]$$

and from Eqs. (28) and (29) we observe that $S(k)$ is extensive, doubling the numbers of billionaires, earthquakes or solar flares in the data sets doubles the value of $S(k)$, and it can be seen to be so because the deformation index α' has the precise value to ensure this property. The constraint

$$\sum_{k=0}^{k_{\max}} k p_k^{\alpha'} = \mathcal{K}_{\alpha'} \quad [30]$$

in Eq. (18) for entropy maximization indicates that the phase space, $N_{\min} \leq N \leq N_{\max}$, is highly constrained since the probabilities $p_k < 1$ need to be enhanced up to $p_k^{\alpha'}$, $\alpha' < 1$, in order to obtain a meaningful average of k . In relation to this, notice that $N(k)$ is a monotonously decreasing function with a power law feature. This phase-space contraction is extreme for the case of Zipf law because α' reaches its minimum value of zero. For a system with normal occupation of phase space, the number of configurations grow exponentially and $S(k)$ above becomes extensive in k for index value $\alpha' = 1$ whereas the phase space in the most contracted stage the number of configurations grow only linearly and and this linearity is preserved in $S(k)$ when $\alpha' = 0$.

In Fig. 3 we show the same data in Figs. 1 and 2 but this time plotted in deformed logarithmic scales with deformation indexes $\alpha \simeq 2$ and $\alpha' \simeq 0$. Data in these scales are displayed linearly and should be fitted by the theoretical expressions Eqs. (7) and (24) if these equations represent the behavior of the data.

Discussion

We have shown that size-rank distributions with power-law decay for moderate and large values of rank obey Tsallis statistics. The small-rank behavior that departs from the power law is also well reproduced by the deformed exponential expression in Eq. (7) for $N(k)$. For the specific data we presented (US billionaires, California earthquakes and solar-flare intensities) the values of the exponential deformations were found to be $\alpha \simeq 2$, the value needed to obtain the classical Zipf law. In order to advance further in the characterization of the apparent relationship between rank distributions and generalized statistical mechanics, such as that of Tsallis, we rederived Eq. (7) for $N(k)$ from a maximum entropy procedure. This was done in accordance to the consideration of validity of only the first three Shannon-Kinchin axioms [11, 12]. Under these conditions duality of entropy expressions appears according to the use of two different constraints. In doing this we introduce the (unnormalized) distribution $p_k = 1/N(k)$, actually $N(k)$ is the number of data for the same rank k (playing the role of a partition function) [4, 5]. We obtain equality of the entropy

expressions $S_1[p_k] = S_2[p_k]$ in Eqs. (26) and (27) and a companion rank distribution expression for p_k , Eq. (24). As it is known [11, 12] the two entropies $S_1[p_k]$ and $S_2[p_k]$ correspond to the dual deformation indexes α and $\alpha' = 2 - \alpha$. We have enquired as to the different roles of the two entropy expressions and identify the physically relevant information carried by each one. We found that the value of the index α fixes the distribution's power-law exponent for $N(k)$ and that the dual index $\alpha' = 2 - \alpha$ ensures the extensivity of the deformed entropy. Finally, we argued that the value $\alpha = 2$, that corresponds to the classical Zipf law, manifests as $\alpha' = 0$ that we interpret as a extreme contraction of the phase space from which the data originates.

ACKNOWLEDGMENTS. G.C.Y and A.R. gratefully acknowledge the hospitality of the Santa Fe Institute. Support by DGAPA-UNAM-IN100311 and CONACyT-CB-2011-167978 (Mexican Agencies) is acknowledged. G.C.Y. was supported by the Scientific Research Projects Coordination Unit of Istanbul University with project number 36529. M.G.-M. acknowledges the generous support of Insight Venture Partners and the Bryan J. and June B. Zwan Foundation.

1. See J.G. van der Galien (2003) in <http://en.wikipedia.org/wiki/ZipfsLaw>
2. Schroeder M. (1991) *Fractals, Chaos, Power Laws*, W.H. Freeman and Company.
3. Gell-Mann M. (1994) *The Quark and the Jaguar: Adventures in Simple and Complex*, W.H. Freeman and Company.
4. Altamirano, C.; Robledo, A. (2011) Possible thermodynamic structure underlying the laws of Zipf and Benford *Eur. Phys. J.B.*, 81:345-351.
5. Robledo, A. (2011) Laws of Zipf and Benford, intermittency, and critical fluctuations, *Chinese Sci. Bull.*, 56:3645-3648.
6. Tsallis, C. (2009) *Introduction to Nonextensive Statistical Mechanics: Approaching a Complex World* (Springer: New York, NY).
7. [http://www.forbes.com/billionaires/tab:overall country:United States](http://www.forbes.com/billionaires/tab:overall%20country:United%20States)
8. Southern California Earthquake Data Center, <http://www.data.scec.org/>
9. <http://tuvalu.santafe.edu/~aaronc/powerlaws/data.htm>
10. Pietronero L., Tosatti E., Tosatti V., Vespignani A. (2001) Explaining the uneven distribution of numbers in nature: the laws of Benford and Zipf, *Physica A*, 293:297-304.
11. Hanel, R; Thurner, S; Gell-Mann, M. (2011) Generalized entropies and the transformation group of superstatistics. *Proc Natl Acad Sci USA* 108:63906394.
12. Hanel, R; Thurner, S; Gell-Mann, M. (2012) Generalized entropies and logarithms and their duality relations. *Proc Natl Acad Sci USA* 109:19151-19154.
13. Hanel, R; Thurner, S; Gell-Mann, M. (2014) How multiplicity determines entropy and the derivation of the maximum entropy principle for complex systems. *Proc Natl Acad Sci USA* 111:6905-6910.
14. Abe, S. (2000) Axioms and uniqueness theorem for Tsallis entropy *Phys. Lett A* 271:74-79.
15. Baldovin, F. and Robledo, A. (2004), Nonextensive Pesin identity: Exact renormalization group analytical results for the dynamics at the edge of chaos of the logistic map, *Phys. Rev. E* 69:045202.
16. Robledo, A. (2013) Generalized Statistical Mechanics at the Onset of Chaos, *Entropy*, 15:5178-5222.




Fig. 1. Three examples of ranked data that appear to display power-law behavior along an interval of rank values. The top panels show data for the wealth of billionaires in the US [7]. The middle panels present data for the energy released by earthquakes in California [8]. The bottom panels provide data for the intensity of solar flares [9]. In the left panels the data is shown in logarithmic scales, whereas the right panels show the same data in log-linear scales. See text for description.




Fig. 2. The same three examples in Fig. 1 are fitted with the expressions in Eqs. (7) and (20). As it can be seen in the figure the values of α needed for fitting are close to $\alpha \simeq 2$ and $\alpha' = 2 - \alpha \simeq 0$. The value $\alpha = 2$ gives the classical Zipf law exponent, whereas the value $\alpha' = 0$ indicates extreme configuration-space contraction. See text for description.




Fig. 3. The same three examples in Figs. 1 and 2 plotted in $\ln_{\alpha}(N(k)/N_{max})$ (left) and $\ln_{\alpha'}(p_k/p_{min})$ (right) scales. Data plotted in these scales are designed to display linear behavior if the theoretical expressions in Eqs. (7) and (20) are fulfilled by the data.