

# Mathematics as a Tool?

Juergen Jost

SFI WORKING PAPER: 2016-09-020

SFI Working Papers contain accounts of scientific work of the author(s) and do not necessarily represent the views of the Santa Fe Institute. We accept papers intended for publication in peer-reviewed journals or proceedings volumes, but not papers that have already appeared in print. Except for papers by our external faculty, papers must be based on work done at SFI, inspired by an invited visit to or collaboration at SFI, or funded by an SFI grant.

©NOTICE: This working paper is included by permission of the contributing author(s) as a means to ensure timely distribution of the scholarly and technical work on a non-commercial basis. Copyright and all rights therein are maintained by the author(s). It is understood that all persons copying this information will adhere to the terms and constraints invoked by each author's copyright. These works may be reposted only with the explicit permission of the copyright holder.

[www.santafe.edu](http://www.santafe.edu)



**SANTA FE INSTITUTE**

# Mathematics as a Tool?

## The Role of Mathematics in the Era of Data Science

Jürgen Jost\*

September 13, 2016

### Abstract

I analyze the new emerging role of mathematics for extracting structure from data. This role is different from the traditional role of mathematics as a tool for other sciences that provided a theoretical framework in which experiments acquired meaning, with which the quality of data could be assessed and which could be explored with the formal methods of mathematics. The challenge for mathematics now is to handle data without such theoretical guidance from other disciplines. This calls for new principles that are structural, abstract and independent of specific content.

## 1 The traditional distinction: Pure and applied mathematics

Traditionally, there has been the distinction between pure and applied mathematics. Pure mathematics – so the story goes – discovers, creates and investigates abstract structures (see [51]) for their own sake, while applied mathematics applies existing mathematical tools and develops new ones for specific problems arising in other sciences. The interaction between pure and applied mathematics takes place in both directions. Applied mathematics utilizes concepts and methods developed in pure mathematics, and problems from diverse applications in turn stimulate the development of new mathematical theories. And traditionally, those problems arose within a clear conceptual framework of a particular science, most notably physics. In this essay, I want to argue that this distinction between pure and applied mathematics is no longer useful – if it ever was –, and that the challenge of large and typically rather diverse data sets, typically arising from new technologies instead of theoretically understood

---

\*Max Planck Institute for Mathematics in the Sciences, Inselstr.22, 04103 Leipzig, Germany, and Santa Fe Institute, Santa Fe, NM 87501, USA

and experimentally testable concepts, not only calls for new mathematical tools, but also necessitates a rethinking of the role of mathematics itself.

In fact, already in the past, what was called applied mathematics often developed domain independent methods. And this domain independence typically led to a gain in generality and mathematical depth. Statistics, for instance, has become so powerful precisely because it developed methods and concepts that apply to essentially any field, from particle physics to the social sciences. And the error estimates and convergence rates provided by numerical analysis are valid for any application of a particular numerical method in the engineering sciences or elsewhere. Nevertheless, such applications usually took place within a particular field with a well developed theoretical framework that provided interpretations for the statistical or computational results obtained. And in other cases, like modeling with differential equations, the specific properties of a concrete scientific theory yielded key ingredients for the mathematical structures. The development of the underlying scientific theory and the mathematical model often went hand in hand, and they deeply depended upon each other. Mathematical modeling was an essential ingredient of the scientific strategy, and in that sense, mathematics was more than a tool. In contrast, nowadays mathematicians are often confronted with data sets of obscure quality, perhaps even of dubious origin, and without any firm theoretical foundation. Even the distinction between meaningful data and meaningless noise may not be clear at all.

Therefore, as data collection these days is typically ahead of theoretical understanding, mathematics should radically face the lack of theory and look at what there is, the data, and see what it can do with them. And what I want to call for are not ad hoc methods for every concrete data set, but rather an abstract analysis of the deeper structural challenges. Of course, modern mathematics is developed and sophisticated enough to provide appropriate and helpful tools for basically any data set, but this by itself is too narrow a scientific perspective. For me as a mathematician, mathematics is more than data analysis. We need a conceptual rethinking.

## 2 What is the role of mathematics in the sciences?

Let us take a look at the situation from the perspective of the sciences. It seems to me that in the presence of data, there are two different, but interwoven issues:

1. The epistemic question, or the role of theory: Can we have data without a theory? And if not, does the theory have to be specific for the domain from which the data are collected?
2. The ontological question, or the role of models: Do we need, or have to postulate, specific objects underlying the data? What is it that the data tell us something about, and how can we, or should we model that?

This distinction may sound a little like issues debated between scientific realists, positivists, and empiricists. But even if we adopt the latter stance and use, for instance, van Fraassen’s criterion that a “theory is empirically adequate if it has some model such that all appearances are isomorphic to empirical substructures of that model” where “appearances” are “structures which can be described in experimental and measurement reports” ([86], p.64), the problem of data without such an empirically adequate theory remains.

Of course, there are some easy answers that seem rather obvious:

- Data that cannot be interpreted within some theoretical framework are meaningless. Much of current data collection is solely driven by particular technologies rather than by scientific questions. As Sir Peter Medawar put it, “No new principle will declare itself from below a heap of facts”.<sup>1</sup> And for “facts”, we might want to substitute “data”.
- If we simply analyze data with intrinsic formal tools whose precise functioning we may not even understand, then science becomes agnostic [64]. We need to know first about what the data are telling us something. That is, we need an underlying structure from which the phenomena revealed by the data are derived. The meaning of data depends on the context from which they are collected.

These concerns are clearly valid. Nevertheless, I think that the issues are somewhat more subtle.

1. To what extent does theory have to be domain specific? Or more precisely, what is the relationship between general theoretical issues – to be elaborated below – and domain specific ones? Or more positively, will the current situation of “big data” lead to new types of theories that start from a data intrinsic rather than a domain specific perspective, and could that possibly lead to theoretical insights at a higher level of abstraction?
2. What is the right balance between a purely phenomenological approach and one that starts with a model involving underlying objects? Such objects could be either the carriers of the properties revealed by the data or at least offer formal structures to which quantities measured on the data are isomorphic.

A traditional perspective would consider mathematics only as a tool when – possibly very large – data sets are to be analyzed. One of the theses of this essay is that this process itself, data analysis, can and already has become an object of mathematical research. Thus, mathematics not only serves as a powerful tool, but by reflecting this role, gains a new perspective and lifts itself to a higher level of abstraction. Then, the domain of such mathematical inquiry no longer is a specific field, like physics, but mathematical technique itself. Mathematics then is no longer, if it ever was, a mere formal tool for science, but becomes the science of formal tools.

---

<sup>1</sup>I learned this quote from Peter Schuster.

Another thesis of this essay is more concerned with the traditional role of mathematics as the formal analysis of models arising from specific domains. I shall argue here against the principle of reductionism. The thesis will be that in the empirical domains where regularities have been identified that could be cast into mathematical structures, the relations between the different levels at which such structures emerge are very intricate. A simple formal structure at one level typically can neither be derived from an even simpler structure at a lower level, nor can be used to identify the relevant structure at a higher level. The structures at both lower and higher levels can be substantially more complex. Again, it is a general task for mathematical research to identify and more deeply understand the principles of such transitions between different scales and levels. That is, what is at issue are not so much the phenomena at the different levels, but rather the transition laws between levels. Much is known here already, both from the side of mathematics and that of physics, but a more abstract and fundamental mathematical theory is still missing.

There might even be unexpected connections between those two aspects, a mathematics of and not only for data analysis on one side, and the principles of transitions between different levels on the other side. At least, there is the question of how such transitions can be inferred from the data. I regard this as a fundamental question for any abstract theory of data.

Before addressing these issues in detail, I should point out that there is another, perhaps even more fundamental and important role for mathematics in the sciences, although that role will not be discussed in this essay. This consists in providing a framework for conceptual thinking and formal reasoning. This is valid across all disciplines, and actually badly needed in many of them. I shall address this aspect elsewhere.

### 3 The role model of physics

In some way or another, there is a particular role model behind many discussions, that of classical physics. According to that perspective, every science should strive towards such a model. We should have objects whose properties and interactions can be formally described by mathematical relations, in particular, differential equations. And we should be able to conduct experiments whose outcomes can be understood within an encompassing theoretical framework. Deviations from that role model are regarded as deficits.

Newton's theory combined the terrestrial dynamics as developed by Galilei and others and Kepler's laws of celestial motions in a unified theory of gravity. The basic objects were pointlike masses. Such a concept would have appeared meaningless to Descartes who considered extension as the basic quality of matter. In Newtonian dynamics, however, the essential feature is the mass of an object, and this allowed him to work with extensionless points carrying a mass as idealized objects. Newton's theory allows for an exact solution of the two-body problem.<sup>2</sup> Besides points, there are other fundamental constituents of physical

---

<sup>2</sup>The one-body problem, however, was eventually understood as being ill posed. As Leibniz

theories. These are the fields, like the electromagnetic one. In physical theories, fields are considered to be no less real than material objects. In particular, they also carry forces which make physical interactions possible. Also, while the two-body problem admits an exact solution in Newton's theory, this is no longer true for the three-body problem. Although that problem is perfectly well posed within Newton's theory of gravity, it can in general no longer be solved in closed form, unless particular symmetries pertain. This was first realized by Poincaré. Nevertheless, one can develop approximation schemes or compute numerical solutions of the differential equations to any desired degree of accuracy with sufficiently large computer power. At a more theoretical level, the stability against perturbations becomes a subtle issue, as developed in KAM theory (after Kolmogorov, Arnol'd, and Moser), see [81, 84]. Also, there are mathematical examples [89, 74] where internal fluctuations in such a system can grow without bounds until one of the bodies is ejected to infinity.

The paradigmatic objects of this theory are the celestial bodies in the solar system, and the theory is concerned with their orbits. Even when the theory is restricted to the basic configuration of the sun, the earth, and the moon, there are more than two such objects. And these objects are by no means pointlike, but rather extended and possessing a non-homogeneous internal structure. If one models them in more detail, the resulting theory becomes much more complicated, and at best one may hope for numerical approximations, but these will inevitably be of limited scope and validity.

But this then raises the question why such a simple theory as Newton's is applicable at all to such a complicated configuration, and in particular, why numerical solutions of Newton's equations lead to such accurate descriptions and predictions of the movements of the planets. (For the sake of the argument, we ignore here the corrections necessitated by Einstein's theory.) When we change the scale, either going down to the internal structure of the celestial bodies, or up to configurations of many gravitating objects, no such simple theory applies.

Is Newton's theory then a completely singular instance, and would we therefore be ill-advised to consider it as a role model of a scientific theory? Well, it is not completely singular, and in order to understand the issues involved better, let us consider another example.

The fundamental equation of quantum mechanics is Schrödinger's equation

$$\sqrt{-1}\hbar\frac{\partial\phi(x,t)}{\partial t} = -\frac{\hbar^2}{2m}\Delta\phi(x,t) + V(x)\phi(x,t) \quad (1)$$

for the quantum mechanical state  $\phi(x,t)$  at position  $x$  and time  $t$ , where  $\Delta$  is the Laplace operator, a second order partial differential operator,  $V(x)$  is the potential at  $x$ , and  $\hbar, m$  are physical constants. The Schrödinger equation no longer describes the deterministic behavior of pointlike objects, but rather

---

first saw, any physical theory has to be concerned with relations between objects, and an irreducible point mass can only entertain relations with other such objects, but not with itself, because there is no fixed external frame of reference independent of objects. The latter aspect is fundamental in Einstein's theory of general relativity. (See for instance the exposition by the author in [72].)

the evolution of probabilities. The dynamics of these probabilities are still deterministic. And in fact, for the hydrogen atom, the Schrödinger equation is exactly solvable. For more complicated atoms, we encounter the difficult numerical problems of quantum chemistry. The reason why it applies so well to the hydrogen atom is that its nucleus consists of a single proton which under normal conditions essentially behaves like an elementary, indivisible (“atomic” in the literal sense) particle. In fact, however, the proton is not elementary, but is composed of more elementary particles, the quarks. That is, the fact that the hydrogen atom can be described by the Schrödinger equation does not have a theoretical, but an empirical reason, the so-called quark confinement. Of course, one may then try to derive this from the standard model of particle physics, but that is a different issue. (For the theoretical background, see e.g. [87].) What remains is the empirical validity of Schrödinger’s equation. The standard model, in contrast, is a model whose deeper justification still is a matter of intense debate. There are approaches like string theory (see e.g. [31, 70, 48, 47, 91]) which, however, are beyond the range of experimental testing for the time being. Why Schrödinger’s equation applies so well (at least to the hydrogen atom) then remains mysterious at a deeper level. And it even applies to other atoms, although it is no longer exactly solvable already for the Helium atom. This is analogous to Newton’s equation and the three-body problem.

The equations of Newton and Schrödinger may possess certain universal properties, and some aspects might be considered with the modern tools of renormalization groups, see for instance [4], but the question why such equations appear at particular scales and not at others remains. Thus, my question is not why a mathematical description of nature is possible, or why mathematics is so effective as a tool for describing physical reality, but rather why it works so well at particular scales or under certain circumstances, but not in other situations. Historically, when the mechanical natural philosophy of the 17th century could capture so well the movements of celestial bodies, explain the acceleration of falling bodies or the parabolic trajectories of canon balls or identify the conservation laws of inelastic collisions, it was expected that this success could easily be extended to other domains and achieve, for instance, a mechanical explanation of animals, and, as some, like La Mettrie, believed, even of humans. Of course, this scientific program turned out to be a miserable failure, and biology as a modern science got off the ground only in the 19th century, on the basis of completely different principles than those of the mechanical natural philosophy of the 17th century.

I shall return in Section 4 to the difference between the theoretical validity of an equation and its exact solvability, as this is important for our topic. Before that, however, I’ll proceed to larger physical scales. In the area of the physics of nano- and microstructures, on the one hand, there are atomic or molecular models, and on the other hand, there are continuum models. Either type of model can yield accurate descriptions within its range of validity and make precise predictions. One might then argue that the discrete (atomic or molecular) models are the basic ones that capture the underlying objects whereas the

continuum models are purely phenomenological. This, however, will miss the fundamental mathematical question, the relationship between these two types of models and the transition between them when changing the scale. Again, this is an aspect to which I need to return. For instance, it seems that in many respects, the Navier-Stokes equations correctly – whatever that means – or at least adequately describe the behavior of fluids. These are continuum equations. The numerical solution, that is, an approximation by a numerical scheme, is one of the most important topics of scientific computation, and by now, there exist numerical schemes that can solve them to a very high degree of accuracy. Nevertheless, within the mathematical theory of partial differential equations, the general existence of a solution to date has not been established, and in fact, this is considered to be a challenging and difficult problem. The issue of turbulence which is still not understood reveals that there are some deep problems lurking here. Interestingly, while turbulent flow seems to be rather chaotic, there might exist some statistical regularities. In any case, the Navier-Stokes equation serve as the most important model in fluid dynamics.

When we move further up with regard to the physical scale, we come to the example already discussed, the dynamics of the orbits of the planets in the solar system which (except for relativistic corrections) are so well described by Newtonian mechanics. Planets, however, are no elementary bodies, but composite objects. So, let us repeat the question why such complicated objects as planets, just think of the geology of the earth, follow such elementary laws. If we go to a smaller scale, everything breaks down, or at least becomes substantially more complicated. For instance, which mathematical or physical theory can explain the rings of Saturn?

And if we go to configurations of large numbers of gravitating bodies, Newton's theory, although applicable in principle (modulo relativistic corrections, again), becomes useless. We should need to change to a statistical description à la Boltzmann [7]. Such a statistical description would still be mathematical; it would derive its power from ignoring the details of the lower level structure.

Physics then loses some of its special role. It seems to be a general phenomenon that in many domains particular scales exist at which a rather simple mathematical model can capture the essential aspects, sometimes even accurately in quantitative terms, while that need no longer hold for smaller or larger scales. It might seem that this occurs less often in domains other than physics, but there do exist positive examples. In chemistry, we have the reaction kinetics. For instance, reaction kinetics of Michaelis-Menten type are important in biochemistry. They depend on important simplifications. In particular, they assume that the chemical substances participating in the process in question are uniformly distributed across the cell. The internal spatial structure of the cell is completely ignored. As another example, the differential equations of Hodgkin-Huxley type model [41] the generation of the spike of a neuron in a quantitatively exact manner, see e.g. [63, 57]. The model operates at the cellular level and describes the dynamics there in a deterministic manner. If one goes down to the lower, molecular, scale, things get much more complicated, and the deterministic quantities in the Hodgkin-Huxley equations become stochastic, that is,

probabilities for the opening and closing of certain ion channels. Similarly, if one moves up and considers interacting systems of neurons, things become quite complicated, and completely different types of models are called for. Thus, we have a relatively simple, formally closed, and quantitatively accurate model at the cellular level, but no such model at the smaller molecular or the larger tissue scale. I shall present the mathematical structure of biochemical reaction kinetics and the biophysics of the Hodgkin-Huxley model in detail below – and readers not interested in those mathematical or physical details can skip those –, but let me first emphasize again the general point. Not only in physical, but also in biological systems, there may exist one or more particular scales at which a gross simplification can lead to simple, but nevertheless quantitatively accurate models, and these scales then correspond to particularly useful levels of description. Typically, however, this is no longer so when we move to either smaller or larger scales. This leads to the question why and how such particular levels emerge, at which such simple quantitative models work. And also, why are such levels apparently so rare?

Let us now describe the mechanism of biological reaction kinetics in some detail to see what is involved. General references for biochemical kinetics are [63], [56], and we follow here the presentation in [49]. The basis is the law of mass action which states that the reaction rate of a chemical reaction is proportional to the concentrations of the reactants raised to the number in which they enter the reaction. That expression is proportional to the probability that the reactants encounter and react with each other. Let us consider the simple reaction



that converts  $S_1 + S_2$  into  $2P$  with forward rate  $k_+$  and backward rate  $k_-$ . That is, when a molecule of substance  $S_1$  encounters one of  $S_2$ , they react and form two copies of  $P$  with rate  $k_+$ . Conversely, two  $P$ -molecules together can decay into a copy of  $S_1$  and a copy of  $S_2$  at the rate  $k_-$ . Thus, the chemical reaction can take place in either direction, but at possibly different rates. If we denote the respective concentrations by  $s_1, s_2, p$  and consider them as functions of time, then (2) leads to the differential equations

$$\begin{aligned} \frac{ds_1}{dt} &= -k_+s_1s_2 + k_-p^2 \\ \frac{ds_2}{dt} &= -k_+s_1s_2 + k_-p^2 \\ \frac{dp}{dt} &= 2(k_+s_1s_2 - k_-p^2) \end{aligned} \tag{3}$$

whose solutions  $s_1(t), s_2(t), p(t)$  then give the respective concentrations at time  $t$  as functions of their initial values. In enzymatic reactions, there also is the complex  $ES$  of the enzyme  $E$  and the substrate  $S$ . The Michaelis-Menten theory makes the simplifying assumption of a quasi-steady state for the complex  $ES$ , that is, its concentration is not changing in time, and this concentration can then be simply computed. This assumption, that the concentration of an

intermediate product remains constant, reduces a multidimensional process to a single equation. The resulting simple systems of ordinary differential equations capture the concentration level of substances involved in biochemical reactions in the cell well enough for many purposes. In contrast to the Schrödinger equation which considers the state  $\phi(x, t)$  as a function of the position  $x$  and time  $t$ , the chemical concentrations  $s_1, s_2, p$  in (3) are considered as functions of  $t$  only. That is, the model assumes that they are homogeneously distributed in the cell. This means that the concentrations are assumed to be the same across the cell at each fixed time  $t$ . This is, of course, a gross simplification, and the model can be refined by allowing for varying concentrations and diffusion effects. The point I want to make here, however, is that even under this (and the further Michaelis-Menten type) simplification, the solutions of systems like (3) can effectively describe the values of concentrations of chemical reactants in cells.

We next turn to the other example discussed above, the Hodgkin-Huxley model [41] for the generation of spikes in neurons,<sup>3</sup> and give a brief description of the model and its dynamics, see e.g. [49, 50] for more details. Formally, the model consists of four coupled (partial) differential equations, but they are more difficult than those for biochemical reactions, and so, we refrain from writing them down here. Instead, we only discuss the qualitative features of the model. The basic dynamic variable is the membrane potential  $V$  of the neuron. The potential is caused by the different densities of charged ions in- and outside the cell. The boundary of the cell, the cell membrane, is impermeable to most charged ions, except for channels that are selectively permeable for certain specific ions. That permeability in turn depends on the membrane potential as well as on the concentration of certain intracellular and extracellular substances. The differential equations of Hodgkin and Huxley then link the temporal changes of these quantities, that is, the membrane potential and three gating variables that control the permeability of the cell membrane for specific electrically charged ions. Concerning the evolution of the membrane potential, the idea is that the time derivative of this potential is proportional to the derivative of the electrical charge, hence to the current flowing. That current in turn is the sum of an internal membrane current and an external current. The latter represents the external input to the cell. The internal membrane current is a sum of specific terms, each proportional to the difference between the potential and some specific rest term. These proportionality factors then depend on the corresponding gating variables. Thus, the dynamical interplay between the membrane current and the gating variables is the key point. When the system is near its resting value and some positive current is injected that lifts the membrane potential  $V$  above some threshold, then a positive feedback between  $V$  and the fast one,  $m$ , among the gating variables sets in. That is, the potential  $V$  rises, and positively charged sodium ions flow in, rising the potential further. Soon, the neuron emits a spike. But then, the two slower gating variables,  $h$  and  $n$ , take over, causing

---

<sup>3</sup>The original model was developed only for a particular type of neuron, the giant squid axon, but similarly models have subsequently been developed for other classes of neurons as well.

an inactivation of the inflow of the sodium ions and reversing the potential by an outflow of potassium ions. The potential then drops even below its resting value, but after some time, during which no further spike is possible, recovers to that latter value, and the neuron is receptive again to repeat the process in response to some new external current. The interaction of two time scales is important for these dynamical properties. The positive feedback between the fast variables  $V$  and  $m$  triggers the spike, whereas the slow variable  $h$  ultimately stops the inflow of positive ions, and  $n$  causes other positive ions to flow outwards, to reverse the effect and bring the potential back to (or, more precisely, below) its resting value. As already mentioned, this particular model was conceived for the giant squid axon. For this neuron, quantitative measurements were easier than for other neurons, and therefore, Hodgkin and Huxley could carefully fit the parameters of their model. Following the pioneering work of Hodgkin and Huxley, then also models for other classes of neurons were developed. While the details are different, the essential principles are the same. Thus, we have quantitative accurate models of the biophysics of neurons that operate at the cellular level, even though the details at the smaller, molecular level are much more intricate. No such simple models exist at that scale.

## 4 Validity vs. computability

We return to Poincaré's insight into the impossibility of an exact solution (a solution in closed form, that is, in the form of an explicit formula for the positions of the bodies involved at every instance of time) of the three-body problem. Even if a mathematical theory of nature were exactly valid in principle – let us ignore the quarks and the internal structure of the planets for the sake of the argument here and assume that for instance, Newton's or Schrödinger's equations were exactly valid –, it could not offer an exactly solvable description of the dynamical behavior of its objects. It is a mathematical question when and under which particular conditions an exact solution by an explicit formula is possible. This has nothing to do with the range of validity of the theory in question. Almost all differential equations cannot be solved by an explicit formula (they do not constitute completely integrable systems in the language of classical mathematics). The search for closed solutions, that is, to demonstrate the complete integrability of a dynamical systems, was one of the highlights of 19th century mathematics. In the 20th century, in the wake of Hilbert, mathematics turned to a different approach. Instead of attempting to construct an explicit solution of a specific differential equation, the problem was converted into showing the existence of solutions for large classes of differential equations. This was the guiding theme in particular for partial differential equations, see for instance [44]. A key point was to separate the abstract question of the existence from the explicit representation of a solution. Once the existence of a solution had been demonstrated from abstract principles, it then became the task of mathematics to understand the properties of such abstract solutions, and in particular to find out under which conditions singularities can be avoided, and

then to derive approximation schemes for such solutions and to convert them into algorithms for their numerical construction with a precise error control. This is basic for all the modern applications of partial differential equations in the engineering sciences and elsewhere. This does not mean, however, that mathematics becomes purely instrumental.

A similar issue arises for the weather forecast. Again, it was Poincaré who first realized the specific aspects of chaotic dynamics. Some meteorological models like the Lorenz equations indeed exhibit chaotic behavior. Therefore, we cannot accurately predict next week's weather, because of the amplifications of tiny fluctuations by chaotic dynamics. It is claimed, however, that the global climate 50 years from now can be predicted within certain bounds, for any scenario of greenhouse gas emissions. The reason why this is feasible is that on a longer time scale, the daily weather fluctuations average out, and only the long term trend remains, and that is precisely what the climate models try to capture. In any case, the peculiarities of chaotic dynamics as exhibited by weather models are not an artefact caused by any deficiency of a model, but are – according to deep mathematical insights – rather grounded in specific structural features of the pertinent equations. The mathematics of chaos goes much beyond the rather simple exponential divergence of individual trajectories, and gains positive insight via such concepts as invariant measures (for details, see e.g. [45]). It can even elucidate the universal nature of many aspects of chaotic dynamics, thereby going much beyond any specific object domain.

In this section, I have described two different trends that distinguish 20th century from 19th century mathematics. On the one hand, the ideal of writing down an explicit solution has been abandoned for the less ambitious aim of approximating a solution to any specified degree of accuracy, or in practice at least as accurately as the computing facilities permit. When one has a good theoretical control of the numerical scheme, then higher accuracy simply requires more computer power. In that regard, mathematics has become more instrumental, concentrating on the computational tools rather than on the explicit formulae. On the other hand, general structural insights into chaotic dynamics teach us that gains in accuracy may require an exponential increase in computational effort and therefore quickly become unrealistic. Thus, in most problems we cannot hope to be able to write down an explicit solution of the mathematical model, and even a very accurate approximation of a solution may not be computationally feasible.

## 5 What are the objects?

As described at length, the objects of Newton's theory are modelled as points that possess masses, but no extension. In solid state physics, physicists are working with objects that have more internal properties and structure and therefore might appear more realistic. Electromagnetic and other fields are objects of physical theories with less material substrate. The nature of subatomic particles is not so clear, even though one might grant them some reality because they

produce discernible effects [35, 25]. When we go to string theory, particles are nothing but excitation modes of completely virtual “objects”, the strings (see e.g. [31, 70, 48, 47, 91]). Such an excitation mode is similar to a Fourier coefficient. There is nothing material about it, and it is a good question in which sense that should be considered as a physical object. Clearly, however, it is a theoretical object, even though it might not possess any independent reality.

Many biological structures are modelled as discrete entities, even though they might be composed of atoms, molecules, cells, or individuals in a complex manner. Mendel’s laws are a good example. Mendel conceived of genes as abstract entities, without any clue about their physical implementation or realization. Although Mendel’s laws are not exactly valid, for many purposes this does not matter. This leads to the question how such discrete entities can emerge from some continuum at a lower scale. Conversely, large ensembles of such discrete objects are often best modelled by continuum models, as in population genetics, see e.g. [42].

In any case, even though it has been discovered in molecular biology that the material substrates of genes are nucleotide sequences, the modern biological conception of a gene is more abstract than such a nucleotide sequence. We refer to [76] and the discussion in the journal *Theory in Biosciences* about this issue [77]. Similarly, the other fundamental biological concept, the species, is more abstract than the collection of individuals composing a population (see for instance [8]). Thus, “objects” in modern biology are abstracta like genes or species that do not directly correspond to physical entities.

This is not fundamentally different in the social sciences, even though there are fewer examples of successful mathematical models. At present, the validity of the mathematical models of macroeconomic theory is somewhat controversial. Likewise, it is an issue of debate to what extent game theory can adequately capture human behavior (see for instance the discussion in [54]). On the other hand, in recent years there has been definite progress in modelling traffic dynamics with quasi-physical theories inspired by microscopic particle physics, gas kinetics or fluid dynamics (see [40]), notwithstanding the fact that individual drivers can behave very differently from each other.

Another issue that has been debated already 200 years ago by social scientists and mathematicians is to what extent the law of large numbers provides not only an empirical, but also a conceptual basis for mathematical modelling of social phenomena and dynamics, see [34]. When the samples are large enough, statistical laws can lead to arbitrarily precise predictions. This is the foundation of such domains as demography or demoscropy. When we recall Boltzmann’s description of statistical ensembles or the above scale transitions, this appears no longer fundamentally different from what has been discussed above for physics and biology.

## 6 Scales and levels

In the preceding, I have already discussed the fact that the difficulty and feasibility of a description or a model of a complex system can be rather different at different levels. In this section, I want to analyze this issue more systematically, drawing upon the collaboration with Nihat Ay, Nils Bertschinger, Robin Lamarche-Perrin, Ekehard Olbrich and Oliver Pfante within the EU Project MatheMACS (Mathematics of Multilevel Anticipatory Complex Systems). Let me start with the terminological distinction between *levels* and *scales*. According to the definitions that we propose

- Scales refer to observables. Scales are determined by the measurement process, the data and their representations at different resolutions. For instance, concerning the dimension of length, we might consider the scales of micrometers, millimeters, meters, and kilometers, or other, smaller or larger ones.<sup>4</sup>
- Levels refer to descriptions or models. Levels arise in the modeling process through the identification of entities that lend themselves to useful analysis. Thus, we could model a biological system at the molecular, cellular, tissue or organismal level.

Clearly, the two concepts are interdependent. Levels rely on measurements taken at characteristic scales. They also rely on a choice of observables that are measured. Scales can be chosen relatively arbitrarily, as long as the relevant measurements can be performed, but a useful choice of scale should depend on the identification of a level. Levels, however, should be distinguished by characteristic properties of the model they allow. That is, at a given level, particular regularities should arise that do not pertain at other levels.

First of all, there are the systems that do not possess any characteristic scale. Such systems have been objects of research in both mathematics and physics. One of the concepts proposed here is that of self-similarity. A self-similar structure looks the same at any scale. Such structures were popularized by Mandelbrot [61] under the name of fractals. The relevant mathematical theory had already been created earlier, by Hausdorff [38] in 1918. Hausdorff had developed a general concept of dimension, and many fractals possess a non-integer Hausdorff dimension. In particular, fractals are in some sense highly irregular. Nevertheless, there also exist such fractals with an integer dimension, and so, mathematically, a more subtle mathematical characterization is necessary, see [82] and the references therein. The concept of a fractal also comes up in the theory of chaotic dynamics where the attractors and the boundaries of their domains of attraction could have such a fractal structure. The corresponding discoveries about iterations of polynomial maps were also made in 1918/19,

---

<sup>4</sup>In different fields, it may be different what is considered as a larger or a smaller scale. In geography, for instance, larger scale means higher resolution, that is, a smaller reduction factor. In other areas, like physics, a larger scale means the opposite. We shall follow the latter terminology. Thus, at a larger scale, many details from a smaller scale may disappear whereas larger structures might become visible.

by Julia and Fatou, see [22] for a more recent description. In the physics literature, scalefree structures are usually characterized by a power-law behavior, as opposed to an exponential decay of correlations. That is, we essentially find correlations at any scale. It turns out that such power-law behavior is rather ubiquitous, from the financial data analyzed by econophysicists to the degree sequences of empirical networks in different domains. From a more theoretical perspective, such systems can even serve as some kind of universal models, when arising as limits of renormalization group flows (see e.g. [11]), and they are basic constituents of conformal field theory and the theory of critical phenomena, see [23, 90].

Of course, no real system can be self-similar at all scales or possess correlations of all orders. Such properties can only hold within a certain range of scales. Nevertheless, as asymptotic idealizations, systems with such properties can be quite useful as theoretical tools.<sup>5</sup> Scalefreeness, however, is not an aspect I want to focus upon here. Let me rather return to a system with well-defined objects at some particular scales. For instance, take a couple of Newtonian particles, at some scale, for instance gas molecules or celestial bodies. When there are only a few of them, their kinetic or dynamical interactions can be described by a mechanical model. For the celestial bodies, that is, when gravity is the active physical force, we have already seen above that for more than two particles, an explicit solution of the equations of motion in general cannot be achieved. But even worse, when the numbers of particles becomes large, like the stars in a galaxy or the gas molecules in a container, the description in terms of classical mechanics is no longer feasible, and we need to turn to a less explicit statistical description à la Boltzmann. Here, a particular configuration of, say, gas molecules, given in terms of their positions and momenta, is a microstate, but at the macrolevel, one only has collective observables like entropy or temperature. These observables are not meaningful at the level of the individual particles, because they represent statistical averages. More precisely, any macrostate can be the result of many different microstates, and the more microstates underlie a given macrostate, the higher the latter's entropy, that is, the more likely it is to occur. (There are some subtle issues here concerning the interpretation of the probabilities involved, see [43], but we do not enter into those here.) In summary, here a deterministic description at the level of individual particles becomes unfeasible and useless due to their large numbers and the impossibility of accurately measuring all their positions and momenta, and instead a statistical description at a higher level is used. It can also be the other way around, that a statistical description at a lower level yields to a deterministic description at a higher level in terms of statistical averages. As fluctuations may average out, these averages themselves may well obey deterministic laws. We see this, of course, in the transition from the quantum level to that of micro- or mesoscopic physics, but for instance also in cells when we go from the molecular to the cellular level. Recall our discussion of the Hodgkin-Huxley equations in Section

---

<sup>5</sup>Batterman [4] and Lenhard[59] emphasize the fact that such idealized systems that arise as asymptotic limits in some theory are employed as models for empirical systems.

3.

Besides the dichotomy between deterministic and stochastic models, there also is the dichotomy between discrete and continuous models. When we move up or down the scales, or better, from one level to the next, again the relation can go either way. Discrete particles (atoms, molecules, stars, cells, individuals, ...) can give rise to continuum models at a higher scale. However, out of underlying continua, also discrete structures can emerge. Partly, these are simply physical processes, for instance when interstellar dust condenses into a star by the force of gravity. Partly, the discrete entities are crucial constructs of models. The gene as a concept in molecular or evolutionary biology emerges from some underlying level of molecular interactions or reproductions of individuals in a population, see [53]. The continuous dynamics of the Hodgkin-Huxley model gives rise to the discrete event of a spike. Or putting it somewhat differently, the neuron as a biological system transforms an analogous input into a binary output, that is, it chooses between the alternatives of spike vs. no spike. Thus, models of information transmission in neural systems can operate with discrete events as the carriers of discrete bits of information. From a formal point of view, such phenomena are genuinely nonlinear, and there exist theoretical concepts, like those of critical threshold, bifurcation, or phase transition, to analyze them, see for instance [45, 49]. Here, however, rather than analyzing the dynamical mechanisms that give rise to such discrete events in a continuous setting, I want to utilize this as an example of a higher level description in discrete terms with an underlying continuous dynamics at a lower level. Interestingly, at the level of discrete events, the spiking pattern of a neuron is often modelled as a Poisson process (see for instance [20, 29, 50]), that is, as a stochastic process, instead of a deterministic one. Here, the reason why one switches from a deterministic to a stochastic description is somewhat different from the case of the Boltzmann gas. In the latter case, a statistical description is called for because of the larger number of particles involved and the infeasibility or impossibility to measure all of them. Here, in contrast, we have a single event, the generation of a spike by a neuron that at one level is modelled by a deterministic dynamical system.<sup>6</sup> Also, this system does not exhibit chaotic behavior that would make long term predictions impossible. Of course, near the critical threshold above which the neuron spikes and below which it returns to rest, the dynamical behavior naturally is unstable in the sense that arbitrarily small perturbations or fluctuations can induce a switch from one state to the other. At another, more abstract, level it is instead modelled as a stochastic process. This is not only simpler, because it dispenses us of having to deal with the subtle nonlinear behavior of the Hodgkin-Huxley system, but it also offers the advantage that we can now look at the relation between the strength of the neuron's input and the single parameter that characterizes the stochastic process, the firing rate of the neuron. Put simply, the hypothesis would be that the neuron fires more frequently on average when it receives a stronger input. (Whether, or perhaps

---

<sup>6</sup>Interestingly, stochastic perturbations of this deterministic system produce genuinely nonlinear effects, see [33, 85].

more precisely, to what extent neuronal systems really employ a rate coding as the preceding might suggest is an unresolved and intensely debated issue in the neurosciences, but here, I do not enter that discussion, and rather refer to [32, 88] for the information theoretical analysis of neuronal spike trains.)

I now turn to the formal analysis of the relation between different levels, following [68]. (Let me also mention earlier results in [79, 30] and many other papers, and the case study in [69].)

We consider a process

$$\phi : X \rightarrow X'. \quad (4)$$

This could, for instance, be the transition from the state of a system at time  $t$  to its state at time  $t + 1$ . We assume that this is a Markov process, in the sense that knowledge of the state at prior times  $t - 1, t - 2, \dots$  does not contain any information beyond that contained in the state at time  $t$  relevant for the state at time  $t + 1$ . That is, the future is conditionally independent of the past given the present. We also have an operator

$$\pi : X \rightarrow \hat{X} \quad (5)$$

that is considered as a projection, coarse graining, averaging, or lumping. This simply means that the  $\hat{\phantom{x}}$  indicates a description at a higher level, with less detail and resolution. The question then is whether we can transform the transition directly at the higher level, that is, whether there exists a process

$$\psi : \hat{X} \rightarrow \hat{X}'. \quad (6)$$

that is self-contained in the sense that it does not depend on anything in  $X$  that is not already present in the upper level  $\hat{X}$ . In other words, we ask whether we can close up the following diagram (make it commutative in mathematical terminology [51]).<sup>7</sup>

$$\begin{array}{ccc} & \psi & \\ \hat{X} & \xrightarrow{\psi} & \hat{X}' \\ \pi \downarrow & & \downarrow \pi' \\ X & \xrightarrow{\phi} & X' \end{array} \quad (7)$$

Expressed as a formula, commutativity means that

$$\psi(\pi(x)) = \pi'(\phi(x)) \quad (8)$$

for all  $x \in X$ .

There exist several criteria to make this precise.

**I Informational closure:** All we need to know to determine the state  $\hat{X}'$  or to predict its statistical properties is already contained in the state  $\hat{X}$ .

The higher process is informationally closed, i.e. there is no information

---

<sup>7</sup>A perhaps somewhat technical point concerning the representation by this diagram: Often, one thinks of a projection as going down, instead of up, and one would then represent  $X$  in the top and  $\hat{X}$  in the bottom row. Since, however, we think of  $\hat{X}$  as a higher, more abstract, level, we rather represent that higher level in the top row of our diagram.

flow from the lower to the higher level. Knowledge of the microstate will not improve predictions of the macrostate. The upper level process is self-consistent in the sense that it does not need to perpetually draw information from or about the lower-level states.

II **Observational commutativity**: It makes no difference whether we perform the aggregation first, and then observe the upper process, or we observe the process on the microstate level, and then lump together the states. In that sense, the upper level process seems autonomous, as once initialized, it appears to unfold on its own, without needing any further updating by details from the lower level process.

III **Commutativity** in the sense of (8): There exists a transition kernel  $\psi$  such that the diagram (7) commutes.

IV **Markovianity**:  $\hat{X}, \hat{X}'$  forms again a Markov process. We recall that we assume that  $X, X'$  yield a Markov process, that is, the current state of lower level process contains everything needed to compute its next state, and it does not need to draw upon past states any further, as these have transmitted all relevant information to the current state. But it does not follow in general that the upper process is Markovian as well, and so, IV indeed is a nontrivial condition. It could compensate a lack of information about the current lower level state by structural constraints in order to also utilize information from past states. In particular, at the upper level, information about past states can improve its prediction about the next state, whereas such information is not useful at the lower level. See the example in [69], and also the discussion in Section 10.

We show in [68] that I implies both II and IV, and II implies III, whereas IV does not imply III in general. Also, for a deterministic process, but not necessarily for a stochastic one, conversely III implies II which in turn implies I. III of course is a formalization of I and II, and as we show, in the deterministic case, they are equivalent. In the stochastic case, the notions turn out to be somewhat different, as information concerns statistical properties, and these are different from the results of particular observations. IV, of course, is a condition that is meaningful only for stochastic dynamics.

## 7 Data without underlying objects?

When we start with data, then an object behind these data is first a hypothesis, a construct. A well discussed example are the gigantic data sets created by modern particle accelerators where the elementary particles that are supposed to produce those data are postulated by theory, but where the only evidence for them has to be extracted from those data (for a conceptual discussion, see [25]). More generally, physics studies phenomena and data, like the scattering matrix in quantum mechanics that records the input and output of a quantum mechanical system. Physicists do not think in terms of objects, in contrast

to the picture of Newtonian physics depicted above. In fact, precisely these scientists are those that drive the data analysis approach also in other domains. For instance, neoclassical economic theory has developed elaborate models for describing the functioning of an (idealized) economy, but then came the so-called econophysicists that simply took the data from financial and other markets and searched for statistical regularities [62]. They used methods from nonlinear time series analysis, for instance. Those methods are not sensitive to the origin of the data, nor do they depend on any object oriented models. Rather, they look for intrinsic regularities in possibly chaotic dynamics [55]. For instance, they identify (chaotic) attractors and determine their (possibly fractal) dimension. These, however, are not ad hoc methods; they are based on deep insights from the theory of dynamical systems. For example, the center manifold principle, see e.g. [45], (which, for instance, is the basis of Haken's slaving principle [36]) tells us that in a dynamical system, typically most directions are quickly relaxing to their equilibrium, and the essential aspects of the dynamics are determined by very few slow variables.

In fact, it had already argued by Koopmans [58] in his review of a book by Burns and Mitchell [9] that measurements of economic quantities without underlying economic concepts are useless; and that time, the criticism was directed against a use of econometric techniques without a theoretical framework grounded in economics, but the same type of criticism would also, and perhaps even more forcefully, apply to the approach taken by econophysics.

We should also remember that Newton's theory that can derive and predict the motion of celestial objects is a rather exceptional feat even in the history of astronomy. Astronomy started with data collection, and these data then were typically organized by models with an ontological status that was dubious at best. The old Babylonians simply interpolated between observational data by assuming that the positions of the celestial bodies varied linearly between the extrema and then suddenly changed direction, see the systematic studies of Neugebauer [65, 66], who speaks of zigzag functions to describe this interpolation scheme. The Ptolemaic model as described in the *Almagest* employed more and more epicycles without really accounting for their ontological status. The Aristotelian system of the celestial spheres was more explicit in that regard, but more difficult to reconcile with the astronomical data. (In fact, Ptolemy also not only tried to represent data as in the *Almagest*, but also developed a planetary hypothesis in which Aristotle's celestial spheres were fattened to provide space for the epicycles needed in Ptolemy's system to account for the astronomical data.) And Kepler had to labor painstakingly for many years through the data compiled by Brahe. He first tried a preconceived scheme, an organization of the solar system in terms of Platonic solids, which he then abandoned because it did not fit the data. In particular, he was concerned with the orbit of Mars. He tried many variants before he finally arrived at fitting the orbit as an ellipse. Thus, by trying to account for the data, in the end he succeeded in discovering his laws of planetary motion. As already discussed, the laws that Kepler had empirically discovered were then derived by Newton from his law of gravity, that is, within an explicit physical model. Of course, without knowing Kepler's

laws, Newton might not have found his theory of gravity.

I should concede, however, that this picture of large data sets that are first collected without an adequate theory and only subsequently inspire deep physical theories is not without exceptions. At least one exception springs to mind, Einstein's theory of general relativity. This theory was developed on the basis of an abstract principle, general covariance, and was only subsequently tested against and confirmed by empirical data.

Also, the traditional view of physics is that it does not collect more or less arbitrary data, but conducts specific experiments whose results acquire their meaning within an established theoretical framework.

On the other hand, however, we should also discuss the approach of Alexander von Humboldt. The novelty of his expeditions rests in the systematic and comprehensive collection of all kinds of data with all the measuring devices available at his time, at a time when there was still very little theoretical understanding of the processes shaping the ecology of the earth. His expedition in South America took some years, but the evaluation of the data collected during that expedition took him several decades. His work then launched the scientific discipline of physical geography and helped and inspired several other scientific fields, even though no single coherent and encompassing theory emerged from his data. Thus, Humboldt had gathered huge amounts of data, but these data did reveal only very few coherent patterns, like the dependence of climatic zones on altitude and latitude. In other words, not only was the model missing, but also the data analysis largely failed in discovering general structures.

## 8 Towards an abstract theory

In fact, it is a quite general finding that there exist profound analogies between models in physically very different domains. It is actually one of the main driving forces of mathematics to consider the corresponding structures and relations abstractly and independently of any particular instantiation and to work out general theories. This will then make it possible, in turn, to apply these mathematical theories to new domains. It is then irrelevant whether such a domain constitutes some independent reality or whether one simply has a collection of data. For instance, similar statistical phenomena show up in quantum mechanics, in the analysis of biological high-throughput data, or in the description of social phenomena.

Understanding the laws and regularities of the transition between scales (see e.g. [67] for the mathematical background), or in a more ambitious formulation, a mathematical approach towards the issue of emergence (for instance, [52]), is a theoretical challenge that transcends individual domains. Nevertheless, it should lead to fundamental insight into the structure of reality, at an abstract level.

In particular, it is a fundamental question in the theory of complex systems to what extent general laws apply across different domains and disciplines, and where the applicability of a general theory ends and a more concrete modelling

of the details of the specific system is required. For instance, which analytical concepts and mathematical tools apply simultaneously to cells, neural systems, psychological systems, and societies, or at least to several of them, and where do the specific peculiarities of each of these fields enter?

In a different direction, we may ask for a theory of structure in high-dimensional spaces. As will be explained below, even though the data may possess a large number of degrees of freedom, these degrees of freedom typically do not vary completely independently in large data sets, but rather obey some nonlinear constraints. Thus, we are dealing with intrinsically lower dimensional geometric structures in high dimensional spaces. Since the particular structure will usually not be known before having analyzed the data, we need to consider spaces of such structures. This leads to new challenges for the mathematical field of geometry. These spaces will then carry a probability density, telling us about the a-priori likelihood of finding them in a particular data set. In statistical terms, we then have some prior hypothesis about the data set, and such a prior then has to be turned into a posterior by Bayes' rule on the basis of the data observed. Of course, this then should be done in an iterative manner. In a different direction, we may employ methods from algebraic topology or metric geometry in order to discover robust qualitative features of specific data sets, see [12, 3].

## 9 The challenge of big data

Both the abundance of huge data sets in almost all disciplines and the availability of the computational power to formally analyze them are quite recent developments. We have the high-throughput data in molecular and cellular biology generated by gene sequencing, microarrays or various spectroscopic techniques, the imaging data in the neurosciences, the email or movement data of mobile phone users, the data of transactions in financial markets at millisecond resolution, the linking patterns of the world wide web, and so on.

And almost uniformly across disciplines, we see a major transition from model driven to data driven approaches. The latter approaches typically depend on the statistics of large data sets. These statistics are utilized automatically, and statistical regularities within large data sets are only used in an implicit fashion, without making them explicit. Examples abound. In molecular biology, the problem of protein folding had for a long time been approached by explicit physical models. The problem consists in predicting the three-dimensional folding pattern from the linear sequence of the amino acids constituting a polypeptide on the basis of molecular attractions and repulsions. As the energy landscape is quite complicated, there are typically many metastable states, and finding the configuration of minimal energy, i.e., the configuration supposedly assumed by the polypeptide in the cell, therefore is computationally quite difficult, even though powerful Monte Carlo type schemes have been developed. Recently, however, it turns out that the best predictions are achieved by data bank searches without any physical model in the background. One simply compares the se-

quence at hand with those sequences where the folding pattern is already known, for instance by X-ray crystallography, and then makes a statistical prediction based on sequence similarities.

Perhaps the preceding becomes clearer when we consider a hypothetical approach to weather forecast. A model driven approach develops a detailed dynamical model of cloud formation and movement, ocean currents, droplet formation in clouds, and so on, and measures the many parameters of such a model, or perhaps also tries to fit some of them on the basis of observations. Such a model would be given by a system of coupled partial differential equations (PDEs), for which one has to solve an initial value problem. The initial values are determined by current measurements on a grid of measurement stations that is dense as possible, naturally with a higher density on land than on the oceans. Since the dynamics described by the PDE model tends to have chaotic aspects (see also the discussion in Section 4), the precision of the measurements of the initial values is of utmost importance in order to have somewhat accurate predictions for a few days. Likewise, the details of the model and its parameters are crucial. In contrast, a data driven approach would like depend on accurate measurements, but it would then try to identify those constellations in the past whose values are closest to those presently recorded, and then use the known weather dynamics from the past for those constellations to predict the weather derived from the current values. This would simply require large data bases of the past weather recordings, and perhaps some scheme of taking weighted averages over similar constellations in the past, but it would not need any model of weather dynamics. Such an approach should then naturally be expected to improve as the data base grows over time.

In the geosciences, the standard scheme consists now in estimating 2-point correlators from noise correlations at different locations (see e.g. [73, 28]), instead of using physical models of, for instance, wave propagation in geological media. The noise sources can be physical or caused by human activities. While the differences matter at a technical level because of different noise characteristics, this does not affect the principle of the method.

Or to present another example that plays a prominent role in [64], microarrays record the simultaneous expression of many different genes in a particular cell condition by exposing the corresponding RNAs in the cell simultaneously to an array of pieces of complementary DNA sequences and then simply recording which of those DNA sequence pieces find RNA partners. The underlying biological rationale is the following. The DNA pieces are part of the genome that is selectively transcribed into RNA which may then be further translated into polypeptides, the building blocks of proteins. Proteins carry out most of the essential operations in a cell. Therefore, mechanisms of gene regulation should ensure that precisely those proteins are manufactured that are needed by the cell in a particular situation. Therefore, precisely those pieces of DNA should be transcribed into RNA that encode the right proteins (there are some problems with this oversimplified account, see [76], but the microarray technology happily ignores them). Thus, a microarray tests the expression patterns and intensities of many DNA segments – which stand for genes in this simplified model – si-

multaneously, because the transcribed RNAs bind to the complementary DNA pieces offered by the microarray.

In computer linguistics, for the purposes of automatic translation, models grounded in syntactic and semantic theory are replaced by statistical techniques that simply utilize cooccurrence patterns of words in large corpora. In particular, Google uses so-called  $n$ -gram models for the purpose of automatic translation. This simply means that one derives the relative frequencies of strings of  $n$  words from databases containing trillions of entries.  $n = 5$  is a typical value. That is, the meaning of a word – insofar as one should still speak about meaning here, in the absence of any semantic concepts – is determined by the environment, that is, a few words preceding and following it, in which it occurs. We shall analyze the conceptual shifts that this implies in more detail in Section 10.

In economics, as already mentioned, techniques from statistical data analysis, like nonlinear time series analysis, are applied to financial data sets in order to find subtle patterns that do not follow from theoretical models. This is the new field called econophysics. Recently, econophysicists also started to develop economic models. They thereby depart from a purely statistical approach that solely analyzes economic or financial data and now build models of economies or financial markets themselves. Their models, however, are in contrast to the classical economic models that employ a so-called representative agent. The latter stands for models with many identical typical agents that are ideally analytically tractable. The agent based models of econophysicists instead utilize very simple, but possible diverse and heterogeneous agents that fit well to large scale computer simulations. That is, instead of a single type of agent that might be rather sophisticated and grounded in economic theory, here many simple agents are employed. These agents depend on a couple of parameters, and the parameter values can and typically do differ between the agents. The representative agents of economics are usually assumed to be fully rational – the argument being that non-optimal agents are quickly exploited by their more clever competitors and thereby driven out of the market. The diverse agents of agent based models are not at all assumed to be rational. They rather follow relatively simple empirical rules, and the purpose of the models is to uncover the collective effects of such behavior of many agents through systematic computer simulations. The validity of the simulation results usually remains unclear, however. An important issue is that the models of (neo)classical economic theory strive for the ideal of exactly solvable equations, or at least for describing the economy by a small and carefully specified set of explicit equations. On the basis of these equations and their solutions, they want to achieve analytical insights into the working of the economy. The agent based models of econophysicists, in contrast, employ much less rigid models, possibly with many parameters, and large numbers of equations that can only be numerically solved. Analytical insight no longer is the foremost aim, and one rather wants to identify unexpected nonlinear effects and critical transitions between different regimes triggered by small variations of crucial parameters.

Even in elementary particle physics, after the confirmation of the Higgs boson, that is, a prediction made by the standard model, at the LHC, in the future

one will probably move towards the automatic analysis of large scale scattering data in order to find unpredicted events that may not fit into any current theoretical model. In any case, already for finding evidence for the Higgs boson, to a large extent automatic data analysis methods have been employed. Such methods may find patterns of correlations or untypical events in huge data sets by completely implicit methods. As described and analyzed for instance in [25], the empirical evidence is gathered in four steps. At the basis, there are position measurements. Adjacent position measurements are then combined into tracks. Adjacent tracks in turn are combined into events. Finally, statistical ensembles of scattering events contain resonances. All this is done in a completely automated way. The tools have no special affinity to particle physics, even though particle physicists are among the people making the most advanced use of them. For instance, so-called neural networks encode the patterns of their training sets in a rather indirect and completely implicit manner in synaptic connection weights (see e.g. [6]). When they are subsequently applied to the real data, they produce corresponding associations which may then be interpreted as patterns in the data set. Neural networks and other such schemes find applications in a wide range of data domains. In fact, the methods themselves also change, and for instance, neural networks are sometimes replaced by other methods like support vector machines.<sup>8</sup>

This phenomenon, the transition from model to data driven approaches, occurs not only in individual and specific domains, but also at the formal level. The field of scientific computing currently undergoes a similar transition. The field of machine learning is concerned with the development of techniques for the automatic analysis of large data sets. Statistics, the science of finding structure in data, is moving into a similar direction, and in fact a profound convergence between machine learning and statistics seems to take place. In particular, the computer intensive Bayesian methods take over much of more traditional parametric statistics. Often, such methods are combined with stochastic search algorithms like Monte Carlo methods or Markov type models. Lenhard[59] also points out the interdependence between data dynamics and computational modelling.

Big data are said to be characterized by large or even huge values of the three Vs, that is,

- **Volume:** often petabytes ( $1000^5 = 10^{15}$  bytes)/day, possibly more
- **Variety:** heterogeneity of data types, representation, and semantic interpretation
- **Velocity:** arrival rate and reaction time

Processing big data requires adapted strategies and methods, and it can be decomposed into five phases

1. Acquisition and recording: filtering, compression, metadata generation

---

<sup>8</sup>Support vector machines are efficient classifiers that use a high-dimensional linear feature space (see [18, 78, 83]).

2. Information extraction, cleaning, and annotation
3. Integration, aggregation, and representation; data base design
4. Analysis and modeling; querying and mining the data
5. Interpretation; visualization; possibilities of interaction of human observers with machine processing

For our purposes, phase 4 is the most relevant. Although computer power is rapidly growing, and cloud computing or even access to supercomputers becomes ever more available, large data sets still may give rise to the “curse of dimensionality”. This means that computer time will increase exponentially with the number of degrees of freedom and therefore quickly exceed even the capacities of supercomputers, unless clever use of specific structures within a data set is made. Therefore, corresponding data analysis techniques are being developed. An example is compressed sensing, see [24, 10, 26]. Here, the idea is to use a specific sparsity assumption, for instance that a large and complicated acoustic data set might be generated by a small number of sound sources only. For instance, the sound sources might be humans carrying on conversations in a crowd. Formally, sparsity means that a high-dimensional vector or matrix possesses only few entries that are different from 0. One does not know beforehand, however, which are the nontrivial ones and how many of them there really are. Thus, there is a not very explicit, but still rather powerful and constraining structural assumption. Or, in a similar vein, one might assume that the data points one has in some high-dimensional space are in fact constrained to some low-dimensional manifold ([5]). This low-dimensional manifold, however, need not be a linear space, and therefore, even though it possesses only few intrinsic degrees of freedom, it may still stretch into many of the ambient dimensions. Thus, the data are supposed to possess some rather constraining type of regularity, but again, this regularity is not explicit, and it is the task of a machine learning scheme to use such an abstract assumption in an efficient manner. When one incorporates such an assumption, one may drastically reduce the number of computations needed to identify the underlying structures from the data set, for instance the sound sources in the above example. The mathematical techniques required are typically nonlinear and therefore on the one hand more flexible in discovering and utilizing hidden regularities and patterns in the data, but then, on the other hand, require more specific, but ideally still automatic, adaptations to the data set at hand.

In many respects, human cognition is still superior to automatic data analysis techniques. In order to utilize the power of human cognition, however, it is necessary to convert the data into a format that is familiar to humans. This leads us into the domain of visualization, a newly emerging scientific discipline between computer science, mathematics, and psychology. Thus, the aim is to develop formal methods that can convert a data set into a form in which humans can easily discern patterns. At a more advanced level, the combination of machine learning tools and human perception may become interactive, which was

the goal of the EU funded research project CEEDs (The Collective Experience of Empathic Data Systems).

## 10 Big data and automatic translation, or a paradigm shift from linguistic theory to language processing

Modern linguistic theory and philosophy is founded upon basic oppositions, for instance between

- *langue* (the abstract system of a language) vs. *parole* (the concrete utterance) (de Saussure [75])
- *diachronous* (across time) vs. *synchronous* (simultaneous) (de Saussure [75])
- *competence* (the ability for the correct syntax of one's native language) vs. *performance* (the actual production of utterances) (Chomsky [13])
- *deep structure* vs. *surface structure*, i.e., a language independent representation in an abstract structure is transformed into a sentence according to the syntax of a specific language (Chomsky [13]), and this transformation obeys the general rules of abstract grammar; the latter have become more general and abstract themselves in the course of Chomsky's work [14, 15]
- *spoken vs written* language, again from de Saussure [75] and emphasized more recently by Derrida [21].

Perhaps the most important opposition is that between de Saussure's [75]

- *paradigmatic* alternatives and *syntagmatic* series; this means that a word at a given position in a sentence can be chosen paradigmatically from a list of words that could grammatically occupy this position, but has to obey the syntactic rules of the sentence. This is also an opposition between *absence* and *presence*, or between *selection* and *constraint*; the alternatives that have not been chosen are absent, but the syntactic constraints are present through the other words in the sentence.

These oppositions are (more or less) formal, but not directly mathematical. They gave rise to a formal theory of language, and for some time, one attempted to utilize that theory for purposes of automatic translation. The idea was essentially to automatically infer the grammatical relationships within a given sentence, that is, the dependencies between the different words and grammatical tokens and the internal references, like the referents of pronouns and anaphora. That grammatical structure could then be transformed into the corresponding structure of another language, and the meanings of the individual words could be correlated with the help of good lexica. For instance, for some time [71] was

popular as a suitable grammatical theory for such purposes. All such attempts, however, have more recently been brushed aside by the big data approach to automatic translation, as developed and pushed in particular by Google. That approach completely ignores, and often even ridicules, linguistic theory, and rather draws upon correlations in huge linguistic corpora. My purpose here is to analyze the underlying conceptual shift and to describe what mathematical structures are behind this approach. Those mathematical structures are completely different from those developed from the context of formal linguistics. And the people working on automatic translation within this paradigm express little interest, if at all, in the conceptual basis of their endeavor. Nevertheless, that will be important for this essay.

Thus, the oppositions sketched above no longer play a role. Instead, we see the *corpus*, that is, a data base of texts in the language(s) in question, as the single basic element. The corpus scales by size. Thus, a small corpus might be seen as corresponding to *parole*, whereas big corpora can approach the *langue* side of de Saussure's dichotomy, and the same applies to Chomsky's dichotomy between *competence* and *performance*. More precisely, there no longer is any such abstract thing as *competence*. The possibly huge collections of linguistic data are all there is. Corpora do not care much whether they have been assembled from contemporary, hence essentially simultaneous texts or whether they result from scanning texts over longer periods of time. The only relevant criteria are of a practical nature, digital availability and computational capacity. Also, corpora can as well be based on automatically recorded spoken language as on written texts. The difference is again largely irrelevant, or at least plays no basic conceptual role. (More precisely, what remains is the technical difference between off-line and on-line processing and translation.)

More importantly and interestingly, the opposition between de Saussure's *paradigmata* and *syntagmata* is also resolved. This best explained through the mathematical concepts of stochastic processes and Shannon information (see e.g. [17, 80, 60, 45]). De Saussure talked about alternative words at a given position in a sentence. This is qualitative, but not quantitative. Information theory, in contrast, would quantify the probabilities of different words to occur at such a position. These probabilities then will not only depend on the abstract properties of that position, but also on the concrete words before and behind it in the sequential string forming the sentence or the text. That is, we do not just have probabilities for the occurrences of words at a given position, but we rather have transition probabilities from a word, or more precisely, a segment of a few words, to the next. In fact, automatic translation today mostly works with pentagrams, that is, overlapping strings of five words.<sup>9</sup> Thus, we no longer have de Saussure's qualitative opposition between selection and constraints, but both aspects are combined and made quantitative within the concept of a stochastic

---

<sup>9</sup>In the terminology of statistical physics, the correlation length of word sequences in texts becomes relatively small after five words. In fact, one would expect that it never becomes exactly zero, that is, there do exist correlations of arbitrary length, whatever small. Thus, in technical terms, when moving from a word to subsequent ones in a text, we have a stochastic process that does not satisfy a Markov property for strings of words of any finite length.

process. (In fact, this is a slight oversimplification. A stochastic process would occur if one has to guess or produce the next word on the basis of those preceding it. Thus, a process describes a temporal sequence as in actual speech. A corpus, however, is not a process unfolding in time, but is simultaneously given as a whole. Therefore, the probabilities are determined not only by the preceding words, but also by the following ones. The underlying mathematical principle, however, is not fundamentally different.) Thus, while the big data approach to automatic translation does not care about an underlying conceptual structure, its analysis nevertheless sheds some light on the limitations of formal language theory, and it points towards mathematical structures that replace the qualitative oppositions by quantitative probabilities. These probabilities are nothing but relative frequencies of pentagrams, i.e., certain word constellations, and they can be automatically computed from the corpora at hand. The larger the corpus, the more accurately such probabilities can be determined, in the sense that further additions to the corpus will likely change them only very little. In a certain sense, this approach asks more precise questions than formal linguistics. When analyzing a temporal sequence, like a spoken text, it would not simply ask at each instant “what could come next?”, but rather “how well can I guess what comes next?”, and the latter is quantified by Shannon’s information [80].

Of course, this approach to automatic translation has its limitations. It cannot capture long range dependencies. In a nutshell, the basic assumption underlying the approach is that the transition between overlapping pentagrams satisfies a Markov property, that is, no information from more than five words back in the sequence is needed for the probability of the next word. From the syntactic perspective, this for instance does not adequately capture the structure of the German language where the different components of the verb can be separated by many words in a sentence. From the semantic perspective, the meaning of some sentence in a text may refer to other, much earlier parts of that text, or even to a context outside the text itself. In formal terms, let us recall also our discussion about Markovianity at different levels in Condition IV in Section 6.

One of the central, but to a large extent still unresolved, issues of linguistics, that between syntax and semantics, between structure and meaning, is simply bypassed by automatic translation. A human translator would first try to extract the meaning of a text and then express that meaning as well as she can in the other language. Automatic translation, in contrast, simply transforms one structure into another one.

## 11 The issue, again and hopefully clearer

We are faced with the following alternative to which no general answer can be given, but which needs to be evaluated in each individual situation.

1. The significance of data collected depends on the specific context from which they are taken, and they cannot be fully understood without that context.

2. Data sets typically possess internal structure, and much of such structure generalizes across different disciplines, domains and contexts, or is at least accessible to context independent methods. Identifying and understanding that structure with formal tools will then in turn enable us to use the data to learn something new and insightful about their context.

This is nothing but the old alternative between specificity and generality, as, for instance, already described by Kant. It is easy to deplore too much of an emphasis on either side and support this by supposedly misguided scientific case studies, and to declare a one-sided practice as agnostic or unscientific. The real challenge consists in finding the appropriate balance between the two aspects in each case. Currently, big data sets offer new opportunities for formal methods and computational models, and this may shift the balance for a while.

Concerning the role of mathematics, this might transcend the alternative between a mathematics of content that establishes and analyzes relations between the concepts of a theory and a purely auxiliary mathematics that is relegated to the role of data handling and preprocessing. What should emerge rather is a mathematics that develops new abstract concepts for analyzing and representing data spaces. For instance, a partial merging of the domains of statistics, high dimensional geometry, information theory and machine learning might take place. Such a mathematics would help to detect structures, as opposed to either formalizing structures proposed and developed by other scientific disciplines or to offering tools for fitting given data into such a structure. This process can only enhance the role and the importance of mathematics, as it becomes liberated from being instrumentalized by conceptualizations that are externally imposed. Instead of accepting models from particular domains, mathematics would itself propose abstract metamodels, like sparsity, smoothness, or symmetry.

## 12 Some consequences

The preceding has important implications for the role of models and hypotheses in the scientific process. From the perspective of Bayesian statistics, we begin the process of scientific inquiry with prior hypotheses constructed by us, in whatever way we may find reasonable or on the basis of whatever prior experience we may have. During the process, such a prior hypothesis gets transformed into a posterior one on the basis of the observations made or the data acquired. This is achieved by applying a fixed rule, that discovered by Bayes. The role of models then is relegated to provide sets of parameters that have to be fitted to the data. This is in stark contrast to the ideal of a model in physics. Such a model should contain only very few, and ideally no free parameters at all that are not theoretically determined but need to be measured. Even for such a nearly ideal model, one may question whether empirical adequacy should constitute a proof of the correctness of the model. In a model with many free parameters that need to be fitted to the data, certainly empirical adequacy can no longer count as a proof of the correctness of the models, and indeed, in complex situations, the term “correct model” may lose its meaning entirely. **There no longer is**

**such a thing as a correct model.** There may only be a superior fit of the parameters to the data collected according to unknown probabilities. See [37] for a more detailed analysis of this issue. Alternatively, we might be seeking regularities in data on the basis of certain structural hypotheses, as described in Section 9. Again, such structural hypotheses like sparsity do not come from the data at hand, but are applied by us in order to get some handle on those data. In a certain sense, they constitute a structural prior, although that prior is usually not updated in a Bayesian manner.

While all this may sound rather agnostic, in practice one might get quite far with those schemes of Bayesian updates, parameter fitting, and structural hypotheses. It is challenge for mathematics to analyze this issue theoretically.

Putting it somewhat differently: Data analysis depends on prior structural assumptions. That could be the prior of a Bayesian approach, or it could be the choice of a family of models as in parametric statistics. It could be a general structural hypothesis like sparsity, a low-dimensional manifold (for instance arising as the center manifold of a dynamical system), certain invariances or symmetries etc. This is the epistemic side. At the ontological side, is there anything underlying the data matching those assumptions? One may argue that these assumptions are our constructions, and that therefore there is no guarantee of any correspondance with the data source. However, a simple theory can sometimes enable us to discover specific phenomena that would otherwise not emerge from beneath the heap of data. One may also argue, as I have tentatively done in this essay, that in some situations, such assumptions could be justified by the structures from which the data are derived, but there is no guarantee for that. The question then is whether those instances where it holds – we have discussed Newton’s theory of planetary motion, the Schrödinger equation, biochemical reaction kinetics, the Hodgkin-Huxley equations or traffic dynamics – are just lucky coincidences, or whether there are more systematic structural aspects of reality – whatever that is – that make our speculations sometimes so successful.

## 13 The dream of mathematics

Does this bring mathematics closer to its dream of a pure structural science, abstract and independent of specific content? We have argued already above that mathematics seeks regularities. Perhaps these regularities are discovered in specific domains, but hopefully, they should apply also in other domains, and ideally, they should be universal. For instance, while Lie’s theory of symmetry groups was originally motivated by the symmetries of the systems of classical mechanics, the theory as such is abstract. In particular, through the work of W.Killing and E.Cartan, it lead to the classification of all possible continuous symmetries (see [39] for the history). This turned out to be of fundamental importance for quantum mechanics, and even further for quantum field theory. It permeates much of pure mathematics. It is also relevant in all applications

where continuous symmetries arise. Even more generally, the notion of a group (Lie groups are a special type of continuous groups), as developed by Gauss and Galois, first arose from studying solutions of algebraic equations, that is, a specific mathematical problem, but it then became something of a paradigm of a mathematical structure as such, and it is now important in almost all fields of mathematics, as well as in many areas of physics and other disciplines.

Perhaps data science enables further steps in this direction. Here, even the original problems are no longer domain specific, as the symmetries and invariances of the systems of classical mechanics, but by their very nature already general and abstract when they apply to all kinds of large data sets. It is then natural that the mathematical tools developed to investigate these problems are at least as abstract as those problems themselves. Thus, data science may lead to an abstract theory of structures, that is, the purest form of mathematics. Incidentally, this brings fields of mathematics into focus that hitherto have been considered as applied and not pure mathematics. Statistics is an example. On the one hand, it can be mathematically treated in terms of the geometry of families of probability distributions, see e.g. [1, 2], and on the other hand, high-dimensional statistics and machine learning might also become a science of distributions of geometric objects in high-dimensional spaces, again independent of any specific content. But there should be mathematical structures even more abstract and general than that.

## Acknowledgements

The present paper is a slightly modified version of a chapter that will appear in a book “Mathematics as a Tool”, edited by Martin Carrier and Johannes Lenhard, to appear in Boston Studies in the Philosophy and History of Science, Springer, and which is the result of a project at the ZIF in Bielefeld, Germany. I am very grateful to my partners in the ZIF project, Philippe Blanchard, Martin Carrier, Andreas Dress, Johannes Lenhard and Michael Röckner, for their insightful comments and helpful suggestions which have contributed to shaping and sharpening the ideas presented here. Martin Carrier and Johannes Lenhard also provided very useful comments on an earlier version of my text. Some of my own work discussed here was supported by the ERC Advanced Grant FP7-267087 and the EU Strep “MatheMACS”.

## References

- [1] S.I.Amari, H.Nagaoka, D.Harada, Methods of information geometry, Amer.Math.Soc., 2007
- [2] N.Ay, J.Jost, H.V.Lê, L.Schwachhöfer, Information geometry, monograph, to appear
- [3] M.Bačák, B.B.Hua, J.Jost, M.Kell, A.Schikorra, A notion of nonpositive curvature for general metric spaces, Diff.Geom.Appl. 38, 22–32, 2015

- [4] R. Batterman, *The Devil in the Details: Asymptotic Reasoning in Explanation, Reduction, and Emergence*. Oxford Studies in Philosophy of Science. Oxford University Press, 2002
- [5] M.Belkin, P.Niyogi, Laplacian Eigenmaps for Dimensionality Reduction and Data Representation, *Neural Comp.* 15, 1373-1396, 2003
- [6] C.Bishop, *Neural networks for pattern recognition*, Oxford Univ.Press, 1995
- [7] L.Boltzmann, *Entropie und Wahrscheinlichkeit*, Ostwalds Klassiker 286, Harri Deutsch, 2000
- [8] O.Breidbach, J.Jost, Working in a multitude of trends. Species balancing populations, *J.Zool.Res.Evol.Syst.*42, 202-207, 2004
- [9] A.Burns, W.Mitchell, *Measuring business cycles*, National Bureau of Economic Research, *Studies of Business Cycles*, No.2, New York, 1946
- [10] E.Candès, J.Romberg, T.Tao, Stable signal recovery from incomplete and inaccurate measurements, *Comm.Pure Appl.Math.*59, 1207-1223, 2006
- [11] J.Cardy, *Scaling and renormalization in statistical physics*, Cambridge Univ.Press, 1996
- [12] G.Carlsson, Topology and data, *Bull.Amer.Math.Soc.* 46, 255-308. 2009
- [13] N.Chomsky, *Aspects of the theory of syntax*, Cambridge, Mass: MIT Press, 1965
- [14] N.Chomsky, *Lectures on Government and Binding*, Dordrecht: Foris, 1981
- [15] N.Chomsky, *The Minimalist Program*, Cambridge, Mass: MIT Press, 1995
- [16] F.Close, *Particle physics. A very short introduction*, Oxford Univ.Press, 2004
- [17] T.Cover and J.Thomas, *Elements of information theory*, Wiley, 1981
- [18] N.Christianini, J.Shawe-Taylor, *An introduction to support vector machines and other kernel-based learning methods*, Cambridge Univ.Press, 2000
- [19] L.Daston, P.Galison, *Objectivity*, Zone Books, 2007
- [20] P.Dayan, L.F.Abbott, *Theoretical neuroscience*, MIT Press, 2001
- [21] J.Derrida, *L'écriture et la différence*, Paris, 1967
- [22] R. L. Devaney, *An introduction to chaotic dynamical systems*. Benjamin/Cummings, Menlo Park 1986

- [23] P.Di Francesco, P.Mathieu, D.Sénéchal, Conformal field theory, Springer, 1997
- [24] D.Donoho, Compressed sensing, IEEE Trans.Inform.Theory 52, 1289-1306, 2006
- [25] B.Falkenburg, Particle metaphysics. A critical account of subatomic reality, Springer, 2007
- [26] M.Fornasier, H.Rauhut, Theoretical foundations and numerical methods for sparse recovery, de Gruyter, 2010
- [27] S.Foucard, H.Rauhut, A mathematical introduction to compressive sensing, Birkhäuser, 2013
- [28] J.Garnier, G.Papanicolaou, Correlation based virtual source imaging in strongly scattering random media, Inverse Problems 28, 075002, 2012
- [29] W.Gerstner, W.Kistler, Spiking neuron models, Cambridge Univ.Press, 2002
- [30] O. Görnerup, M. Nilsson Jacobi, A method for finding aggregated representations of linear dynamical systems, Adv.Complex Syst.13, 199-215, 2010
- [31] M.Green, J.Schwarz, E.Witten, Superstring theory, 2 vols., Cambr.Univ.Press, 1995
- [32] S.Grün, S.Rotter, Analysis of parallel spike trains, Springer, 2010
- [33] B.Gutkin, H.Tuckwell, J.Jost, Inhibition of rhythmic neural spiking by noise: the occurrence of a minimum in activity with increasing noise, Naturwissenschaften 96, 1091-1097, 2009
- [34] I.Hacking, The taming of chance, Cambr.Univ.Press, 1990
- [35] I.Hacking, Representing and intervening, Cambr.Univ.Press, 1983
- [36] H.Haken, Synergetics, Springer, <sup>3</sup>1983
- [37] Hasse, J.Lenhard, to appear in: M.Carrier, J.Lenhard (eds.), Mathematics as a tool, Boston Studies in the Philosophy and History of Science, Springer, in press
- [38] F.Hausdorff, Dimension und äußeres Maß. Math. Annalen 79 (1918), 157 - 179
- [39] T.Hawkins, Emergence of the theory of Lie groups. An essay in the history of mathematics 1869–1926, Springer, 2000
- [40] D.Helbing, Traffic and related self-driven many-particle systems, Rev.Mod.Phys.73, 1067-1141, 2001

- [41] A.Hodgkin and A.Huxley, A quantitative description of ion currents and its application to conductance and excitation in nerve membranes, *J.Physiol.*117, 500–544, 1952
- [42] J.Hofrichter, J.Jost, T.D.Tran, Information geometry and population genetics, in preparation
- [43] E.T.Jaynes, Probability theory: The logic of science, Cambridge Univ.Press, 2003
- [44] J.Jost, Partial differential equations, Springer, <sup>3</sup>2013
- [45] J.Jost, Dynamical systems, Springer, 2005
- [46] J.Jost, Riemannian geometry and geometric analysis, Springer, <sup>6</sup>2011
- [47] J.Jost, Geometry and physics, Springer, 2009
- [48] J.Jost, Bosonic strings, Amer.Math.Soc. and Intern. Press, 2001
- [49] J.Jost, Mathematical methods for biology and neurobiology, Springer, 2014
- [50] J.Jost, Mathematical Neurobiology. Concepts, Tools, and Questions, monograph, to appear
- [51] J.Jost, Mathematical concepts, Springer, 2015
- [52] J. Jost, N. Bertschinger, E. Olbrich, Emergence. A dynamical systems approach, *New Ideas in Psych.* 28, 265–273, 2010
- [53] J.Jost, K.Scherrer, A conceptual analysis of genetic coding and regulation in the light of modern molecular biology, manuscript to be submitted
- [54] A.Kabalak, E.Smirnova, J.Jost, Non-cooperative game theory in biology and cooperative reasoning in humans, *Theory in Biosciences* 134, 17–46, 2015
- [55] H.Kantz, T.Schreiber, Nonlinear time series analysis, Cambr.Univ.Press, 1997
- [56] E.Klipp, R.Herwig, A.Kowald, C.Wierling, H.Lehrach, Systems biology in practice, Wiley-VCH, 2005
- [57] C.Koch, Biophysics of computation, Oxford Univ.Press, New York, 1999
- [58] T.Koopmans, Measurement without theory, *Rev.Econ.Statistics* 29, 161–172, 1947
- [59] J.Lenhard, Coal to diamonds, 2013
- [60] D.MacKay, Information theory, inference, and learning algorithms, Cambridge Univ.Press, 2003

- [61] B. B. Mandelbrot, *The Fractal Geometry of Nature*. Freeman, San Francisco 1982
- [62] R. Mantegna, H. E. Stanley, *An Introduction to Econophysics: Correlations and Complexity in Finance*, Cambridge University Press, 1999
- [63] J.Murray, *Mathematical biology*, 2 vols., Springer, <sup>3</sup>2008
- [64] D.Napolitani, M.Panza, D.Struppa, *Agnostic science. Towards a philosophy of data analysis*, *Found.Sci.*16, 1-20, 2011
- [65] O.Neugebauer, *The exact sciences in antiquity*, Dover, <sup>2</sup>1969
- [66] O.Neugebauer, *Problems and methods in Babylonian mathematical astronomy*. Henri Norris Russell Lecture, 1967, *Astronomical J.* 72, 964–972, 1967
- [67] G.Pavliotis, A. Stuart, *Multiscale methods. Averaging and homogenization*, Springer, 2008
- [68] O. Pfante, N. Bertschinger, E. Olbrich, N. Ay, J. Jost, Comparison between different methods of level identification, *Adv. Complex Syst.* 17, 1450007, 2014
- [69] O. Pfante, E. Olbrich, N. Bertschinger, N. Ay, J. Jost, Closure measures and the tent map, *Chaos* 24, 013136, 2014
- [70] J.Polchinski, *String theory*, 2 vols., *Cambr.Univ.Press*, 1998
- [71] C.Pollard and I.Sag, *Head-Driven Phrase Structure Grammar*, Chicago: University of Chicago Press, 1994
- [72] B. Riemann, *Über die Hypothesen, welche der Geometrie zu Grunde liegen*, edited with a historical and mathematical commentary by J. Jost, Springer, 2013; English translation: *On the hypotheses which lie at the bases of geometry*, Birkhäuser, in press
- [73] M.Ritzwoller, F.C.Lin, W.Shen, Ambient noise tomography with a large seismic array, *C. R. Geoscience*, doi:10.1016, 2011
- [74] D.Saari, Z.Xia, Off to infinity in finite time, *Notices AMS* 42, 538–546, 1995
- [75] F. de Saussure, *Cours de linguistique générale*, published by C. Bally, A. Sechehaye and A. Riedlinger, critical edition by T. de Mauro, Payot, 1995
- [76] Scherrer, K. and Jost, J., Gene and genon concept: Coding vs. regulation, *Theory in Biosciences* 126, 65-113, 2007
- [77] Scherrer, K. and Jost, J., Response to commentaries on our paper Gene and genon concept: Coding vs. regulation, *Theory in Biosciences* 128, 171-177, 2009

- [78] B.Schölkopf, A.Smola, Learning with kernels, MIT Press, 2002
- [79] C.Shalizi, C. Moore, What is a macrostate? subjective observations and objective dynamics, Tech. report, 2003, arXiv:cond-mat/0303625v1
- [80] C.Shannon, W.Weaver, The mathematical theory of communication, ed. R. Blahut, B.Hajek, Univ. Illinois Press, 1998
- [81] C.Siegel, J.Moser, Lectures on celestial mechanics, Springer, 1971
- [82] K.Steffen, Hausdorff-Dimension, reguläre Mengen und total irreguläre Mengen, pp. 185-227 in: E.Brieskorn (ed.), Felix Hausdorff zum Gedächtnis, Vieweg, 1996
- [83] I.Steinwart, A.Christmann, Support vector machines, Springer, 2008
- [84] D. Treschev, O. Zubelevich, Introduction to the perturbation theory of Hamiltonian systems, Springer, 2010
- [85] H.Tuckwell, J.Jost, Analysis of inverse stochastic resonance and the long-term firing of Hodgkin-Huxley neurons with Gaussian white noise, Physica A, 391, 5311-5325, 2012
- [86] B. van Fraassen, The scientific image, Oxford, Clarendon Press, 1980
- [87] S.Weinberg, The quantum theory of fields, 3 vols., Cambr.Univ.Press, 1995, 1996, 2000
- [88] M.Wibral, R.Vincente, J.Lizier (eds.), Directed information measures in neuroscience, Springer, 2014
- [89] Z.Xia, The existence of non-collision singularities in Newtonian systems, Ann. Math.135, 411-468, 1992
- [90] J.-J. Zinn-Justin, Quantum field theory and critical phenomena, Oxford Univ.Press, <sup>2</sup>1993
- [91] B.Zwieback, A first course in string theory, Cambr.Univ.Press, <sup>2</sup>2009